

ACCEPTED VERSION

Mathai Varghese and Graeme Wilkin

Fractional quantum numbers, complex orbifolds and noncommutative geometry

Journal of Physics A: Mathematical and Theoretical, 2021; 54(31):314001-1-314001-18

© 2021 IOP Publishing Ltd.

Originally published: <http://dx.doi.org/10.1088/1751-8121/ac0b8c>

This Accepted Manuscript is available for reuse under a [CC BY-NC-ND licence](#) after the 12 month embargo period provided that all the terms of the licence are adhered to.

PERMISSIONS

<https://publishingsupport.iopscience.iop.org/accepted-manuscripts/>

After acceptance, each Named Author of an article to be published/published on a subscription basis may:

Unless otherwise stated, any reference below to an Embargo Period is a reference to a period of 12 months from the Date of Publication.

3. Post the Accepted Manuscript to an institutional repository or subject repository (in both cases ONLY where non-commercial) after the Embargo Period under a [CC BY-NC-ND licence](#), provided that all terms of the licence are adhered to, and any copyright notice and any cover sheet applied by IOP is not deleted or modified. *The above should satisfy the requirements of research funders for 'green open access', such as the Chinese Academy of Sciences, US National Institutes of Health, NASA, NSF, US Department of Energy, NIST, National Research Council of Canada and Austrian Science Fund, to deposit the outputs of research funded by them in a repository.*

4. Post the Accepted Manuscript to an institutional repository or a subject repository (in both cases ONLY where non-commercial) where necessary to comply with the requirements of the HEFCE REF 2021 open access policy. HEFCE's requirements are as follows:

- For articles with a Date of Acceptance between 1 April 2016 and 31 March 2018 inclusive, the Named Authors may make a Closed Deposit of the Accepted Manuscript to the non-commercial repository within three months of the Date of Publication of the article; or
- For articles with a Date of Acceptance on or after 1 April 2018, the Named Authors may make a Closed Deposit of the Accepted Manuscript to the non-commercial repository within three months of the Date of Acceptance of the article.

In both the above cases, after the Embargo Period, the full text of the Accepted Manuscript may be made available on the non-commercial repository for anyone with an internet connection to read and download. After the Embargo Period a [CC BY-NC-ND licence](#) applies to the Accepted Manuscript, in which case it may then only be posted under that CC BY-NC-ND licence provided that all the terms of the licence are adhered to, and any copyright notice and any cover sheet applied by IOP is not deleted or modified.

You may indicate that the CC BY-NC-ND licence applies after the Embargo Period by including the following wording on the Accepted Manuscript, "This Accepted Manuscript is available for reuse under a [CC BY-NC-ND licence](#) after the 12 month embargo period provided that all the terms of the licence are adhered to" (unless there is a cover sheet applied to it by IOP which already states this).

14 September 2022

ACCEPTED MANUSCRIPT

Fractional quantum numbers, complex orbifolds and noncommutative geometry

To cite this article before publication: Mathai Varghese *et al* 2021 *J. Phys. A: Math. Theor.* in press <https://doi.org/10.1088/1751-8121/ac0b8c>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2021 IOP Publishing Ltd.

During the embargo period (the 12 month period from the publication of the Version of Record of this article), the Accepted Manuscript is fully protected by copyright and cannot be reused or reposted elsewhere.

As the Version of Record of this article is going to be / has been published on a subscription basis, this Accepted Manuscript is available for reuse under a CC BY-NC-ND 3.0 licence after the 12 month embargo period.

After the embargo period, everyone is permitted to use copy and redistribute this article for non-commercial purposes only, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by-nc-nd/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions will likely be required. All third party content is fully copyright protected, unless specifically stated otherwise in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

FRACTIONAL QUANTUM NUMBERS, COMPLEX ORBIFOLDS AND NONCOMMUTATIVE GEOMETRY

VARGHESE MATHAI AND GRAEME WILKIN

ABSTRACT. This paper studies the conductance on the universal homology covering space Z of 2D orbifolds in a strong magnetic field, thereby removing the rationality constraint on the magnetic field in earlier works [3, 29, 25] in the literature. We consider a natural Landau Hamiltonian on Z and study its spectrum which we prove consists of a finite number of low-lying isolated points and calculate the von Neumann degree of the associated holomorphic spectral orbundles when the magnetic field B is large, and obtain fractional quantum numbers as the conductance.

CONTENTS

Introduction	1
1. Preliminaries	3
1.1. The maximal abelian cover	3
1.2. Magnetic translations	3
1.3. The magnetic Schrödinger operator	4
1.4. The noncommutative torus	5
1.5. Noncommutative complex torus and holomorphic vector bundles	6
2. Summary of results	6
3. Discrete values of the spectrum of the Laplacian on the maximal abelian cover	8
3.1. An expression for the eigensections	10
3.2. Spectral gaps around the low-lying eigenvalues	11
4. An Index Theorem	14
5. Chern number of E_{μ_q} via a higher index theorem	16
References	17

INTRODUCTION

This paper is motivated by the fundamental result of Avron, Seiler & Zograf [3], in which they compute the *conductance* and *charge transport* associated to a class of quantum systems. These quantum systems roughly describe the dynamics of a Schrödinger particle on a multiply connected two-dimensional surface with no boundary and finite area. The key insight of [3] is that a local families index theorem due to Quillen splits the conductances into two parts (cf. [3, eqn. (14)]). The first part is (up to an integer factor) the canonical symplectic form on the

2010 *Mathematics Subject Classification*. Primary 32L81, Secondary 58J52 58J90 81Q10.

Key words and phrases. fractional quantum numbers, Riemann orbifolds, holomorphic orbundles, orbifold Nahm transform, noncommutative geometry.

space of Aharonov-Bohm fluxes, which is quantized, thus giving a connection to the Integer Quantum Hall effect (IQHE) [31, 2, 4]. The second part is a complete derivative, therefore it does not affect charge transport, however it does affect the conductance as a fluctuation term. These results were extended by Prieto [29], who computed the transport coefficients associated to the eigensections of low lying eigenvalues of the Schrödinger operator. In our previous work [25], we calculated the transport coefficients associated to the eigensections of low lying eigenvalues for any compact complex 2D orbifold where the magnetic field B is a large fraction.

The goal of this paper is to remove the topological constraints on the magnetic field (rationality) in these earlier works [3, 29, 25] in the literature. To achieve this, we instead study the conductance and charge transport on the universal homology orbi-covering space Z of 2D orbifolds in a strong magnetic field. It turns out that there is a projective unitary action of \mathbb{Z}^{2g} on $L^2(Z)$, known as magnetic translations, which commutes with the self-adjoint Hamiltonian

$$H = \frac{\hbar^2}{2m} \left(\nabla_A^* \nabla_A + \frac{R}{6} \right),$$

where $\nabla_A = d + iA$ is a connection on the trivial line bundle on Z with curvature $(\nabla_A)^2 = i\tilde{B}$, and R is the constant scalar curvature of Z in a hyperbolic metric. We also assume that the magnetic field \tilde{B} is a constant multiple of the volume form $\theta d\text{vol}_Z$ for some large value of $\theta \in \mathbb{R}$. In this paper, we do not consider an electric potential, which has to be real valued as we want the Hamiltonian to be self-adjoint, and therefore it will not be compatible with the holomorphic structure that we wish to use. Hence ours is a toy model for the (fractional) quantum Hall effect and more generally a method for producing fractional quantum numbers. Since Z is a noncompact Riemann surface, the Hamiltonian H acting on $L^2(Z)$ typically has continuous spectrum. However, we show in Theorem 2.1 that there are finitely many eigenvalues $\{\mu_1, \dots, \mu_{m-1}\}$ of the Hamiltonian H that are isolated in the spectrum of H and which are near zero. Let $\{E_{\mu_j}, j = 1, \dots, m-1\}$ denote the corresponding eigenspaces. In Theorem 2.1 we show that these eigenspaces consist of holomorphic sections and compute the rank $r_j = \dim_{\mathbb{C}} E_{\mu_j}$. Let $\{P_j, j = 1, \dots, m-1\}$ denote the respective orthogonal projections $P_j : L^2(Z) \rightarrow E_{\mu_j}$ and let τ denote the von Neumann trace. Then by [23], E_{μ_j} has a $\bar{\partial}$ -operator $\bar{\nabla} = P_j(\bar{\partial})^{r_j}$, where $P_j \in M(r_j, A_\theta)$ and A_θ is a complex noncommutative torus in dimension $2g$, generated by the magnetic translations. An open question is whether there is a connection on E_{μ_j} such that $\bar{\nabla}^2 = 0$ that is $\bar{\nabla}$ is a *flat $\bar{\partial}$ operator* in the sense of [23], which is known to be true in the commutative case as shown in [29]. We remark that the Hamiltonian H can be interpreted as a family of Hamiltonians parametrised by A_θ which can be viewed as the noncommutative analog of the Jacobian variety of X .

Hyperbolic 2D surfaces are typically prohibited experimentally as they cannot be isometrically embedded in three dimensional Euclidean space [14]. A recent striking development in [16] indisputably shows that lattices of certain resonators can be used to produce artificial photonic materials in an effectively curved space, including the 2D hyperbolic plane. In

particular, they conducted numerical tight-binding simulations of hyperbolic analogs of the Kagome lattice and demonstrated that they display a flat band, similar to that of their Euclidean counterpart. The authors of [16] also constructed a proof-of-principle experimental device which realizes a finite section of non-interacting heptagon-kagome lattice.

We mention alternate approaches to (fractional) quantum numbers on hyperbolic space. These approaches are for smooth surfaces [8, 7, 9], for orbifolds [17, 18, 19] for the bulk-boundary correspondence [24] and for orbifold symmetric products [20]. These papers use operator algebras and noncommutative geometry methods, in contrast to the holomorphic geometry methods used in this paper. For a recent analysis of the IQHE, see [15], where the generating functional, the adiabatic curvature and the adiabatic phase for the IQHE are studied on a compact Riemann surface with integral magnetic field, but using holomorphic methods inspired by [3].

Acknowledgements. VM thanks the Australian Research Council for support via the Australian Laureate Fellowship FL170100020. He gave a talk partly based on this paper at the conference, *Topological Phases of Interacting Quantum Systems*, BIRS, Oaxaca, Mexico, June 2–7, 2019. GW would also like to thank the University of Adelaide for their hospitality during the development of this paper. His visit was funded by FL170100020.

1. PRELIMINARIES

1.1. The maximal abelian cover. Let X be a compact Riemann surface of genus g with a hyperbolic metric. The first homology is the abelianisation of $G := \pi_1(X)$

$$H_1(X) = G/[G, G].$$

Let $p : Z \rightarrow X$ be the maximal abelian cover with $\pi_1(Z) = [G, G]$. For any $x \in X$, we have $p^{-1}(x) \cong H_1(X)$, and so the commutator subgroup has infinite index.

A theorem of Griffiths [13, 4.2] shows that the commutator subgroup $[G, G]$ is then a free group. The homology $H_1(Z)$ is then the corresponding free abelian group, and hence the cohomology with coefficients in any ring R is the dual $H^1(Z, R) \cong \text{Hom}(H_1(Z), R)$. Putman [27, Lem. A.1] shows that the generators of the commutator subgroup can be realised geometrically as loops on the surface that bound a one-holed torus.

The infinite cover $p : Z \rightarrow X$ defined above is induced from the Abel-Jacobi inclusion $X \hookrightarrow J(X)$ and the universal cover of the Jacobian

$$(1.1) \quad \begin{array}{ccc} Z & \xrightarrow{\tilde{\iota}} & \widetilde{J(X)} \\ \downarrow p & & \downarrow q \\ X & \xrightarrow{\iota} & J(X) \end{array}$$

To see this, let $q : Z' \rightarrow X$ be the covering of X induced from the universal cover of the Jacobian. Note that Z' is connected. Then choose $x \in X$ and note that the action of

$H_1(X) \cong H_1(J(X)) \cong \pi_X(J(X))$ on $p^{-1}(x) \cong H_1(X) \cong q^{-1}(x)$ is the same for both Z and Z' . Therefore the two coverings are isomorphic.

1.2. Magnetic translations. The Abel-Jacobi inclusion $\iota : X \hookrightarrow J(X)$ induces an isomorphism $\iota^* : H^1(J(X), \mathbb{R}) \cong H^1(X, \mathbb{R})$, and the induced map $\iota^* : H^2(J(X), \mathbb{R}) \rightarrow H^2(X, \mathbb{R})$ is surjective. In fact, if the magnetic field is $B = \theta\omega$, where ω is the Kähler 2-form on X and $\theta \in \mathbb{R}$, then one has,

$$[B] = [\theta\omega] = \iota^* \left[\frac{\theta}{g} \Theta_X \right] \in H^{1,1}(X, \mathbb{R})$$

where $[\Theta_X] \in H^{1,1}(J(X), \mathbb{R})$ is the Theta divisor.

Let $B' = \frac{\theta}{g} \Theta_X$ be the closed (1,1)-form on the Jacobian $J(X)$. Then $\tilde{B}' = q^*B'$ is a closed 2-form on $\widetilde{J(X)}$. Since $\widetilde{J(X)}$ is contractible, $H^2(\widetilde{J(X)}, \mathbb{R}) = \{0\}$, so that $\tilde{B}' = dA'$ for some 1-form A' on $\widetilde{J(X)}$.

Now $0 = \gamma^*(\tilde{B}') - \tilde{B}' = d(\gamma^*A' - A')$, so that $\gamma^*A' - A'$ is a closed 1-form on $\widetilde{J(X)}$ for all $\gamma \in \Gamma = H_1(X, \mathbb{Z}) = H_1(J(X), \mathbb{Z})$. Since $\widetilde{J(X)}$ is contractible, $H^1(\widetilde{J(X)}, \mathbb{R}) = \{0\}$, so that $\gamma^*A' - A' = d\phi'_\gamma$, where ϕ'_γ is a smooth function on $\widetilde{J(X)}$ normalised by $\phi'_\gamma(\tilde{\iota}(x_0)) = 0$ for all $\gamma \in \Gamma$ and for some $x_0 \in Z$.

Define $A := \tilde{\iota}^*A' \in \Omega^1(Z)$ and $\phi_\gamma := \tilde{\iota}^*\phi'_\gamma$, and note that $\tilde{B} = p^*B = dA$. Moreover, ϕ_γ satisfies the identity

$$(1.2) \quad \phi_{\gamma_2}(x) + \phi_{\gamma_1}(\gamma_2 \cdot x) - \phi_{\gamma_1 + \gamma_2}(x) = \phi_{\gamma_1}(\gamma_2 \cdot x_0)$$

for all $x \in Z$ and $\gamma_1, \gamma_2 \in \Gamma$. Now define $\sigma : \Gamma \times \Gamma \rightarrow U(1)$ by

$$(1.3) \quad \sigma(\gamma_1, \gamma_2) = e^{i\phi_{\gamma_1}(\gamma_2 \cdot x_0)}.$$

Then the above identity (1.2) shows that σ satisfies the cocycle condition

$$(1.4) \quad \sigma(\gamma_1, \gamma_2)\sigma(\gamma_1 + \gamma_2, \gamma_3) = \sigma(\gamma_1, \gamma_2 + \gamma_3)\sigma(\gamma_2, \gamma_3), \quad \gamma_1, \gamma_2, \gamma_3 \in \Gamma.$$

For each $\gamma \in \Gamma$, define operators on $L^2(Z)$ by

$$\begin{aligned} U_\gamma f(x) &= f(\gamma^{-1}x) \\ S_\gamma f(x) &= e^{i\phi_\gamma(x)} f(x). \end{aligned}$$

Then the above identities imply that $T_\gamma := U_\gamma \circ S_\gamma$ satisfies $T_{\gamma_1}T_{\gamma_2} = \sigma(\gamma_1, \gamma_2)T_{\gamma_1\gamma_2}$, and hence defines a twisted or projective action of Γ on $L^2(Z)$.

1.3. The magnetic Schrödinger operator. Let $\nabla_A = d + iA$ be a connection on the trivial line bundle on Z . Then the curvature of ∇_A is $(\nabla_A)^2 = i\tilde{B}$. Consider the magnetic Schrödinger operator

$$(1.5) \quad H = \frac{\hbar^2}{2m} \left(\nabla_A^* \nabla_A + \frac{R}{6} \right),$$

where R is the constant scalar curvature of Z . Then $U_\gamma \nabla_A = \nabla_{\gamma^{-1}*A} U_\gamma$ and $S_\gamma \nabla_{\gamma^{-1}*A} = \nabla_A S_\gamma$, so that $T_\gamma H = HT_\gamma$ for all $\gamma \in \Gamma$.

Notice that H is bounded below, so the spectral projection

$$P_\lambda = \chi_{(-\infty, \lambda)}(H) \in W^*(\Gamma, \sigma) \otimes B(L^2(\mathcal{F})),$$

where \mathcal{F} is a connected fundamental domain for the action of Γ on Z , $B(L^2(\mathcal{F}))$ denotes the bounded operators on the Hilbert space $L^2(\mathcal{F})$, and $W^*(\Gamma, \sigma)$ denotes the σ -twisted group von Neumann algebra of Γ generated by the magnetic translations acting on $\ell^2(\Gamma)$.

If however λ is in a spectral gap of H , then it is a standard result (cf. [6, Thm. 1]) that the spectral projection belongs to the much smaller algebra,

$$P_\lambda = \chi_{(-\infty, \lambda)}(H) = f(H) \in C^*(\Gamma, \sigma) \otimes \mathcal{K}(L^2(\mathcal{F})),$$

where $\mathcal{K}(L^2(\mathcal{F}))$ denotes the compact operators on the Hilbert space $L^2(\mathcal{F})$, and $C^*(\Gamma, \sigma)$ denotes the σ -twisted group C^* -algebra of Γ and f is a holomorphic function defined in a neighbourhood of $(-\infty, \lambda]$. When \tilde{B} is a constant multiple $\theta d \text{vol}_Z$ of the hyperbolic volume, which is the case that we will focus on in this paper, then $C^*(\Gamma, \sigma)$ is the noncommutative torus A_θ in dimension $2g$.

1.4. The noncommutative torus. Here we recall the definition of the higher dimensional noncommutative torus, its complex structure and K-theory, and the range of the trace and also the 2-trace on K-theory. Let $p = 2g$ and Θ be a $(p \times p)$ skew-symmetric matrix. Then

$$\sigma(\gamma, \gamma') = \exp\left(2\pi\sqrt{-1} \sum_{j < k} \Theta_{jk} \gamma_j \gamma'_k\right), \quad \text{where } \gamma = (\gamma_1, \dots, \gamma_p), \gamma' = (\gamma'_1, \dots, \gamma'_p) \in \mathbb{Z}^p.$$

is a $U(1)$ -valued group 2-cocycle on \mathbb{Z}^p . Let $\mathbb{C}(\mathbb{Z}^p, \sigma)$ denote the twisted group algebra, that is for functions $f_1, f_2: \mathbb{Z}^p \rightarrow \mathbb{C}$ of finite support, the twisted convolution product is

$$f_1 \star_\sigma f_2(\gamma) = \sum_{\gamma_1 + \gamma_2 = \gamma} f_1(\gamma_1) f_2(\gamma_2) \sigma(\gamma_1, \gamma_2)$$

Then $\mathbb{C}(\mathbb{Z}^p, \sigma)$ acts on bounded operators on $\ell^2(\mathbb{Z}^p)$ by the formula above. The operator norm closure $\overline{\mathbb{C}(\mathbb{Z}^p, \sigma)}$ is defined to be the noncommutative torus A_Θ or the twisted group C^* -algebra $C^*(\mathbb{Z}^p, \sigma)$.

There is an abstract definition of A_Θ that is useful to recall. The *noncommutative torus* A_Θ is the universal C^* -algebra with p unitary generators U_j , $1 \leq j \leq p$, subject to the basic commutation relation

$$U_j U_k = e^{2\pi i \Theta_{jk}} U_k U_j.$$

This algebra carries a *gauge action* of the torus \mathbb{T}^p via

$$t \cdot (U_1^{n_1} \dots U_d^{n_d}) = t_1^{n_1} \dots t_d^{n_d} U_1^{n_1} \dots U_d^{n_d}, \quad t = (t_1, \dots, t_d) \in \mathbb{T}^p.$$

There are associated infinitesimal generators δ_j , which are $*$ -derivations, defined by

$$\delta_j(U_k) = 2\pi i \delta_{jk} U_k.$$

The algebra A_Θ also carries a canonical tracial state τ invariant under the gauge action, sending 1 to 1 and sending a monomial $U_1^{n_1} \cdots U_d^{n_p}$ to 0 unless all of the n_j vanish. Because of the commutation relation, any element of A_Θ has a canonical (formal) expansion in terms of the monomials $U_1^{n_1} \cdots U_d^{n_p}$. Since every element of A_Θ has a unique expression $\sum a_{\vec{n}} U_1^{n_1} \cdots U_d^{n_p}$, so

$$\tau \left(\sum_{\vec{n}} a_{\vec{n}} U_1^{n_1} \cdots U_d^{n_p} \right) = a_{\vec{0}}$$

The *smooth noncommutative torus* \mathcal{A}_Θ consists of all elements $\sum a_{\vec{n}} U_1^{n_1} \cdots U_d^{n_p}$ such that the coefficients $a_{\vec{n}}$ form a rapidly decreasing sequence in $\mathcal{S}(\mathbb{Z}^p)$. It is also the smooth vectors for the gauge action of \mathbb{T}^p on A_Θ . It follows that \mathcal{A}_Θ is the domain of the (powers of) derivations δ_j^k that are the infinitesimal generators of the gauge action. There is a natural continuous cyclic 2-cocycle τ_c on \mathcal{A}_Θ . Let $f_0, f_1, f_2 \in \mathcal{A}_\Theta$. Then

$$(1.6) \quad \tau_c(f_0, f_1, f_2) = \sum_{i=1}^g \tau(f_0(\delta_i(f_1)\delta_{i+g}(f_2) - (\delta_{i+g}(f_1)\delta_i(f_2)))).$$

where c is the area 2-cocycle on \mathbb{Z}^p corresponding to the symplectic form.

The inclusion $\mathcal{A}_\Theta \hookrightarrow A_\Theta$ is known to induce an isomorphism in K-theory, $K_\bullet(A_\Theta) \cong K_\bullet(\mathcal{A}_\Theta)$. The range of the trace on K-theory has been computed [5, 11]:

$$(1.7) \quad \tau(K_0(A_\Theta)) = \mathbb{Z} + \sum_{0 < |I| < p} \text{Pf}(\Theta_I)\mathbb{Z} + \text{Pf}(\Theta)\mathbb{Z},$$

where I runs over subsets of $\{1, \dots, p\}$ with an even number of elements, and Θ_I denotes the skew-symmetric submatrix of $\Theta = (\Theta_{ij})$ with $i, j \in I$. The formula (see section 1 in [22])

$$e^{\frac{1}{2}dx^t\Theta dx} = \sum_I \text{Pf}(\Theta_I)dx^I$$

is key to this computation, together with the twisted L²-index theorem [21].

We can also compute the range of a certain higher trace on K-theory. Let Θ_X denote the theta divisor. In real coordinates, $\Theta_X = \sum_{i=1}^g dx_i \wedge dx_{i+g}$. Let c be the group cocycle that corresponds to Θ_X . Then setting $I_i = \{i, i+g\}$, one has

$$(1.8) \quad \tau_c(K_0(\mathcal{A}_\Theta)) = \sum_{i=1}^g \sum_{I_i \subset I} \text{Pf}(\Theta_{I \setminus I_i})\mathbb{Z}$$

where I runs over subsets of $\{1, \dots, p\}$ with an even number of elements, and $\Theta_{I \setminus I_i}$ denotes the skew-symmetric submatrix of $\Theta = (\Theta_{ij})$ with $i, j \in I \setminus I_i$. The method of proof again uses [5, 18] and can be deduced from Corollary 5.7.2 in [26].

1.5. Noncommutative complex torus and holomorphic vector bundles. The following is recalled from [23]. Define a tangent space of \mathcal{A}_Θ to be the (commutative) Lie algebra $\mathfrak{g} = \text{span}(\delta_1, \dots, \delta_{2n})$. A *complex structure* on A_Θ and \mathcal{A}_Θ is a choice of an endomorphism J of \mathfrak{g} satisfying $J^2 = -1$. It thus defines an isomorphism $\mathfrak{g}_\mathbb{C} \cong \mathfrak{g}^{\text{hol}} \oplus \mathfrak{g}^{\text{antihol}}$ as a direct sum of holomorphic and antiholomorphic tangent spaces, namely the $\pm i$ -eigenspaces of J . There

is a similar splitting of the complexified cotangent space $\mathfrak{g}_{\mathbb{C}}^*$. The pair (A_{Θ}, J) will be called a *noncommutative complex torus* of complex dimension n .

A *vector bundle* E over \mathcal{A}_{Θ} will mean a finitely generated projective (right) module. A *holomorphic vector bundle* E will mean such a bundle equipped with a *holomorphic connection* $\bar{\nabla}$, meaning a map $E \rightarrow E \otimes (\mathfrak{g}^{\text{antihol}})^*$ satisfying the Leibniz rule

$$(1.9) \quad \bar{\nabla}_{\bar{\partial}_j}(e \cdot a) = \bar{\nabla}_{\bar{\partial}_j}(e) \cdot a + e \cdot \bar{\partial}_j(a).$$

Note that any vector bundle of rank r can be equipped with a holomorphic connection simply by writing $E = p(\mathcal{A}_{\Theta})^r$ for some projection p , and then defining $\bar{\nabla} = p(\bar{\partial})^r$.

More interesting are *flat holomorphic connections*, which satisfy the flatness condition $(\bar{\nabla})^2 = 0$. It is *not* the case that every vector bundle has a flat holomorphic connection.

2. SUMMARY OF RESULTS

This section contains the main results of the paper.

Theorem 2.1. *Let H be the magnetic Schrödinger operator of (1.5).*

- (1) *Let $m > 0$ be the largest integer such that $\theta - m(2g - 2) > 0$. If $\lambda = \mu_q$ for some integer q such that $0 \leq q < m$, where*

$$\mu_q = (2q + 1)\theta - q(q + 1)(2g - 2),$$

then $\lambda = \mu_q$ is an isolated point in the spectrum of H . It follows that the spectral projection $P_{\lambda} \in A_{\theta} \otimes \mathcal{K}(L^2(\mathcal{F}))$.

- (2) *The eigenspace $E_{\lambda} = \text{Im}(P_{\lambda})$ for λ as above, consists of holomorphic sections.*
- (3) *The von Neumann dimension of E_{λ} , $\dim_{\tau}(E_{\lambda})$, is equal to $\tau(P_{\lambda})$, where $\tau : A_{\theta} \otimes \mathcal{K}(L^2(\mathcal{F})) \rightarrow \mathbb{C}$ is the von Neumann trace. Then E_{μ_q} is infinite dimensional since we show that*

$$\dim_{\tau}(E_{\mu_q}) = (2q + 1)(1 - g) + \theta > 0,$$

and so the von Neumann dimension $\dim_{\tau}(E_{\mu_q})$ of the spectral subspace grows linearly with q .

- (4) *Let $\lambda = \mu_q$. The Chern number of E_{λ} is*

$$(2.1) \quad \tau_2(P_{\lambda}, P_{\lambda}, P_{\lambda}) = \tau(P_{\lambda} dP_{\lambda} dP_{\lambda}) = 2g.$$

In particular, it is an integer and it is independent of q . In the case of an orbifold given as a quotient $X = X'/\Gamma$ of a smooth Riemann surface X' by a finite group Γ , the Chern number is a rational number

$$(2.2) \quad 2g - 2 + \#(R/\Gamma) + \frac{2 - n}{|\Gamma|},$$

where R is the ramification divisor of the ramified cover $X' \rightarrow X$. Here we use the well known fact that the noncommutative torus $C^*(\Gamma, \sigma)$ comes with a canonical cyclic 2-cocycle,

$$\tau_2(f_0, f_1, f_2) = \sum_{j=1}^g \tau(f_0 \delta_j f_1 \delta_{j+g} f_2 - \delta_{j+g} f_1 \delta_j f_2)$$

for $f_k \in \mathbb{C}(\Gamma, \sigma)$. This is derived to be the conductance 2-cocycle τ_K in Corollary 5, [8], see also page 73 in [18].

The subsequent sections contain the proof of these results. Parts (1) and (2) follow from Theorem 3.2, part (3) is proved in Lemma 4.2 and the results of part (4) are contained in (5.1) and (5.2).

3. DISCRETE VALUES OF THE SPECTRUM OF THE LAPLACIAN ON THE MAXIMAL ABELIAN COVER

The main result of this section is Theorem 3.2, proving the first two parts of Theorem 2.1.

Let $\mathcal{E} \rightarrow X$ be a complex line bundle. Since Z is a noncompact surface then the pullback $\tilde{\mathcal{E}}$ is topologically trivial, and any holomorphic structure on $\tilde{\mathcal{E}}$ is also trivial (see for example [12, Thm. 30.4]), however $\tilde{\mathcal{E}}$ admits many different gauge-equivalent structures as a holomorphic bundle, or equivalently as a Hermitian bundle with a unitary connection. In order to normalise the eigenvalues of the Laplacian, we will fix a metric on X of constant Gauss curvature $\chi(X)$ so that $\text{vol}(X) = 2\pi$, and use the pullback metric on the maximal abelian cover $Z \rightarrow X$. Fix a Hermitian metric on $\tilde{\mathcal{E}}$ and a Hermitian connection ∇^θ , and let $\theta = i * F_{\nabla^\theta} \in \mathbb{R}$ be the curvature, which we assume from now on to be constant. In general θ can be any real number; in the special case that ∇^θ is the pullback of a constant curvature connection on a line bundle $\mathcal{E} \rightarrow X$, then the Chern-Weil formula $\text{deg}(\mathcal{E}) = \frac{i}{2\pi} \int_X * F_{\nabla^\theta} = \theta$ shows that $\theta = \text{deg}(\mathcal{E}) \in \mathbb{Z}$.

Elliptic regularity shows that the eigensections of $(\nabla^\theta)^* \nabla^\theta$ are smooth. In the following, we will use the Sobolev space $H^2(Z, \mathcal{E}) \subset L^2(Z, \mathcal{E})$ as the domain of the Laplacian, on which $(\nabla^\theta)^* \nabla^\theta$ is self-adjoint, and continuous as an operator $H^2(Z, \mathcal{E}) \rightarrow L^2(Z, \mathcal{E})$.

The main result of this section is Theorem 3.2, which shows that the low-lying eigenvalues of the Laplacian $(\nabla^\theta)^* \nabla^\theta$ are the discrete values given by (3.4), and that the corresponding eigensections are images of holomorphic sections of the associated bundle $K^{-q} \otimes \tilde{\mathcal{E}}$ given by (3.5).

Decomposing ∇^θ into $(0, 1)$ and $(1, 0)$ parts induces operators ∂^{∇^θ} and $\bar{\partial}^{\nabla^\theta}$, which satisfy the following identities for the Laplacian and the curvature

$$(3.1) \quad (\nabla^\theta)^* \nabla^\theta = \Delta_{\bar{\partial}}^\theta + \Delta_{\partial}^\theta, \quad i * F_{\nabla^\theta} = \Delta_{\bar{\partial}}^\theta - \Delta_{\partial}^\theta.$$

Combining these shows that

$$(3.2) \quad (\nabla^\theta)^* \nabla^\theta = 2\Delta_{\bar{\partial}}^\theta + i * F_{\nabla^\theta} = 2\Delta_{\bar{\partial}}^\theta + \theta,$$

and so the eigensections of $(\nabla^\theta)^*\nabla^\theta$ with eigenvalue μ correspond exactly to eigensections of $\Delta_{\bar{\partial}}^\theta$ with eigenvalue $\frac{1}{2}(\mu - \theta)$. The first consequence of this is that $\mu \geq \theta$, since $\Delta_{\bar{\partial}}^\theta$ is non-negative. Secondly, we see that the sections in the kernel of $\Delta_{\bar{\partial}}^\theta$ (corresponding to the holomorphic sections of \mathcal{E}) are eigensections of $(\nabla^\theta)^*\nabla^\theta$ with eigenvalue θ .

The basic example of such a connection on a trivial bundle with trivial metric on \mathbb{C}^g is

$$\bar{\partial}^{\nabla^\theta} = \bar{\partial} + \frac{1}{4}i\theta z d\bar{z}, \quad \partial^{\nabla^\theta} = \partial - \frac{1}{4}i\bar{z} dz,$$

where we use the shorthand $z d\bar{z} := \sum_{k=1}^g z_k d\bar{z}_k$, $\bar{z} dz := \sum_{k=1}^g \bar{z}_k dz_k$. One can easily check that $e^{-\frac{1}{4}i\theta|z|^2}$ is in $\ker \bar{\partial}^{\nabla^\theta}$ and therefore is an eigenfunction of $(\nabla^\theta)^*\nabla^\theta$ with eigenvalue θ . Since the inclusion $Z \hookrightarrow \mathbb{C}^g$ induced from the Abel-Jacobi map is a holomorphic embedding, then the restriction of $e^{-\frac{1}{4}i\theta|z|^2}$ to $Z \subset \mathbb{C}^g$ is also in $\ker \bar{\partial}^{\nabla^\theta}$ and therefore an eigenfunction of the Laplacian.

Therefore the lowest eigenvalue of $(\nabla^\theta)^*\nabla^\theta$ is determined by the curvature θ , and the corresponding eigensections are determined by the holomorphic sections of \mathcal{E} . The next theorem extends this result to show that the higher eigenvalues of $(\nabla^\theta)^*\nabla^\theta$ also have an explicit description. Most importantly, they are discrete in the interval $[0, \mu_m)$, where m and μ_m are defined below. For compact surfaces, this result is due to Prieto in [30], and the proof below involves extending these techniques to apply to the noncompact infinite genus surface Z .

First, we set some notation. Let T be the tangent bundle of Z , which we assumed to have constant Gauss curvature $\chi(X)$. The bundle T also has a canonical holomorphic structure and Hermitian metric induced from that of Z , and the dual is identified with the canonical bundle K , which then has an induced connection. Define

$$\nabla^{\theta,q} := \tilde{\nabla}^{K^q} \otimes \nabla^\theta$$

to be the connection on $\tilde{\mathcal{E}} \otimes K^q$ for any $q \in \mathbb{Z}$, and decompose into $(1, 0)$ and $(0, 1)$ parts to define the associated operators

$$\partial^{\nabla^{\theta,q}} : \Omega^0(\tilde{\mathcal{E}} \otimes K^q) \rightarrow \Omega^{1,0}(\tilde{\mathcal{E}} \otimes K^q) \cong \Omega^0(\tilde{\mathcal{E}} \otimes K^{q+1})$$

$$\bar{\partial}^{\nabla^{\theta,q}} : \Omega^0(\tilde{\mathcal{E}} \otimes K^q) \rightarrow \Omega^{0,1}(\tilde{\mathcal{E}} \otimes K^q) \cong \Omega^0(\tilde{\mathcal{E}} \otimes K^{q-1}).$$

Remark 3.1. In analogy with (3.1), note that the Laplacians satisfy the identity

$$(3.3) \quad i * F_{\nabla^{\theta,q}} = \Delta_{\partial^{\theta,q}} - \Delta_{\bar{\partial}^{\theta,q}}.$$

In particular, if $\theta - q(2g - 2) > 0$, then $\partial^{\nabla^{\theta,-q}} : \Omega^0(\tilde{\mathcal{E}} \otimes K^{-q}) \rightarrow \Omega^0(\tilde{\mathcal{E}} \otimes K^{-q+1})$ is injective, since $\Delta_{\partial^{\theta,-q}} = \theta - q(2g - 2) + \Delta_{\bar{\partial}^{\theta,-q}}$ is strictly positive.

We then have two sequences of homomorphisms given by composing these operators as q increases or decreases

$$\Omega^0(\tilde{\mathcal{E}} \otimes K^{q-1}) \begin{array}{c} \xrightarrow{\partial^{\nabla^{\theta,q-1}}} \\ \xleftarrow{\bar{\partial}^{\nabla^{\theta,q}}} \end{array} \Omega^0(\tilde{\mathcal{E}} \otimes K^q) \begin{array}{c} \xrightarrow{\partial^{\nabla^{\theta,q}}} \\ \xleftarrow{\bar{\partial}^{\nabla^{\theta,q+1}}} \end{array} \Omega^0(\tilde{\mathcal{E}} \otimes K^{q+1}) \begin{array}{c} \xrightarrow{\partial^{\nabla^{\theta,q+1}}} \\ \xleftarrow{\bar{\partial}^{\nabla^{\theta,q+2}}} \end{array} \Omega^0(\tilde{\mathcal{E}} \otimes K^{q+2})$$

Theorem 3.2. Fix a metric on Z with constant Gauss curvature $\chi(X) = 2 - 2g$, and let m be the largest integer such that $\theta - m(2g - 2) > 0$. For each integer $0 \leq q \leq m$, define

$$(3.4) \quad \mu_q = (2q + 1)\theta - q(q + 1)(2g - 2).$$

Then the spectrum of the Laplacian in the interval $[0, \mu_m)$ consists of the discrete eigenvalues μ_q for each $q = 0, \dots, m - 1$. For each such eigenvalue μ_q , the corresponding space of eigensections E_{μ_q} is equal to

$$(3.5) \quad E_{\mu_q} = \overline{\partial^{\nabla^{\theta, -1}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} (\ker \bar{\partial}^{\nabla^{\theta, -q}})} \subset \Omega^0(\tilde{\mathcal{E}}).$$

In particular, E_{μ_q} and $\ker \bar{\partial}^{\nabla^{\theta, -q}}$ are isomorphic as A_θ modules.

Remark 3.3. These are exactly the same as the eigenvalues in the discrete spectrum of the Laplacian on \mathbb{H}^2 computed by Comtet and Houston [10] (in (3.4) Z has Gauss curvature $-\frac{1}{a^2} = \chi(X)$). This is a nontrivial observation, since the cover $\mathbb{H}^2 \rightarrow Z$ has structure group the infinitely generated free group $\pi_1(Z) = [\pi_1(X), \pi_1(X)]$, and so the L^2 spectrum on Z is not necessarily contained in the L^2 spectrum on \mathbb{H}^2 .

The proof of this theorem is contained in Propositions 3.6 and 3.7 below. Before proving these propositions we first need some basic results about the Laplacian on Z . In the following, the closure will always be taken in the Sobolev space $H^2(Z, \tilde{\mathcal{E}})$, which we use for the domain of the Laplacian.

Lemma 3.4. For each q , the space of sections is a direct sum

$$(3.6) \quad \Omega^0(\tilde{\mathcal{E}} \otimes K^q) \cong \ker \bar{\partial}^{\nabla^{\theta, q}} \oplus \overline{\text{im } \bar{\partial}^{\nabla^{\theta, q-1}}}.$$

The next result describes how the curvature measures the failure of the operators $\partial^{\nabla^{\theta, -(q+1)}}$ and $\bar{\partial}^{\nabla^{\theta, q+1}}$ to commute with the associated Laplacians. The proof is a local calculation as carried out in [30, Prop. 10], and so it also applies to the maximal abelian cover Z .

Lemma 3.5. In the following diagram

$$\begin{array}{ccc} \Omega^0(K^{q+1} \otimes \tilde{\mathcal{E}}) & \xrightarrow{\bar{\partial}^{\nabla^{\theta, q+1}}} & \Omega^0(K^q \otimes \tilde{\mathcal{E}}) \\ \downarrow \Delta_{\bar{\partial}}^{\theta, q+1} & & \downarrow \Delta_{\bar{\partial}}^{\theta, q} \\ \Omega^0(K^{q+1} \otimes \tilde{\mathcal{E}}) & \xrightarrow{\bar{\partial}^{\nabla^{\theta, q+1}}} & \Omega^0(K^q \otimes \tilde{\mathcal{E}}) \end{array}$$

we have

$$(3.7) \quad \Delta_{\bar{\partial}}^{\theta, q} \circ \bar{\partial}^{\nabla^{\theta, q+1}} - \bar{\partial}^{\nabla^{\theta, q+1}} \circ \Delta_{\bar{\partial}}^{\theta, q+1} = -i \bar{\partial}^{\nabla^{\theta, q+1}} * F_{\nabla^{\theta, q+1}}.$$

Equivalently, the above diagram commutes up to a factor of $-i \bar{\partial}^{\nabla^{\theta, q+1}} * F_{\nabla^{\theta, q+1}}$. Similarly, in the following diagram

$$\begin{array}{ccc} \Omega^0(K^{-(q+1)} \otimes \tilde{\mathcal{E}}) & \xrightarrow{\partial^{\nabla^{\theta, -(q+1)}}} & \Omega^0(K^{-q} \otimes \tilde{\mathcal{E}}) \\ \downarrow \Delta_{\bar{\partial}}^{\theta, -(q+1)} & & \downarrow \Delta_{\bar{\partial}}^{\theta, -q} \\ \Omega^0(K^{-(q+1)} \otimes \tilde{\mathcal{E}}) & \xrightarrow{\partial^{\nabla^{\theta, -(q+1)}}} & \Omega^0(K^{-q} \otimes \tilde{\mathcal{E}}) \end{array}$$

we have

$$(3.8) \quad \Delta_{\bar{\partial}}^{\theta, -q} \circ \partial^{\nabla^{\theta, -(q+1)}} - \partial^{\nabla^{\theta, -(q+1)}} \circ \Delta_{\bar{\partial}}^{\theta, -(q+1)} = i\partial^{\nabla^{\theta, -(q+1)}} * F_{\nabla^{\theta, -(q+1)}}.$$

Equivalently, the above diagram commutes up to a factor of $+i\partial^{\nabla^{\theta, -(q+1)}} * F_{\nabla^{\theta, -(q+1)}}$.

3.1. An expression for the eigensections. Recall the definition of m as the largest integer such that $K^{-m} \otimes \tilde{\mathcal{E}}$ has positive curvature, or equivalently such that $\theta - m(2g - 2) > 0$.

The first proposition constructs eigensections of the Laplacian with given eigenvalues.

Proposition 3.6. *For all $0 \leq q \leq m$, the sections in*

$$(3.9) \quad \overline{\partial^{\nabla^{\theta, -1}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} (\ker \bar{\partial}^{\nabla^{\theta, -q}})} \subset \Omega^0(\tilde{\mathcal{E}})$$

are eigensections of $(\nabla^{\theta})^* \nabla^{\theta}$ with eigenvalue

$$\mu_q = (2q + 1)\theta - q(q + 1)(2g - 2).$$

Proof. Let $s_{-q} \in \ker \bar{\partial}^{\nabla^{\theta, -q}} \subset \Omega^0(\tilde{\mathcal{E}} \otimes K^{-q})$, and let $s = \partial^{\nabla^{\theta, -1}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} s_{-q}$. Then Lemma 3.5 implies that

$$(3.10) \quad \begin{aligned} \Delta_{\bar{\partial}}^{\theta} s &= \Delta_{\bar{\partial}}^{\theta} (\partial^{\nabla^{\theta, -1}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} s_{-q}) = \partial^{\nabla^{\theta, -1}} \circ \Delta_{\bar{\partial}}^{\theta, -1} \circ \partial^{\nabla^{\theta, -2}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} s_{-q} \\ &\quad + \partial^{\nabla^{\theta, -1}} (i * F_{\nabla^{\theta, -1}}) \partial^{\nabla^{\theta, -2}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} s_{-q} \\ &= \partial^{\nabla^{\theta, -1}} \circ \Delta_{\bar{\partial}}^{\theta, -1} \circ \partial^{\nabla^{\theta, -2}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} s_{-q} \\ &\quad + (\theta - (2g - 2))s, \end{aligned}$$

since the curvature $i * F_{\nabla^{\theta, -1}}$ of the connection on $\tilde{\mathcal{E}} \otimes K^{-1}$ is constant and equal to $\theta - (2g - 2)$. Repeatedly applying Lemma 3.5 shows that

$$(3.11) \quad \begin{aligned} \Delta_{\bar{\partial}}^{\theta} (\partial^{\nabla^{\theta, -1}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} s_{-q}) &= \partial^{\nabla^{\theta, -1}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} \circ \Delta_{\bar{\partial}}^{\theta, -q} s_{-q} + \left(\sum_{\ell=1}^q (\theta - \ell(2g - 2)) \right) s \\ &= \left(q\theta - \frac{q(q+1)(2g-2)}{2} \right) s, \end{aligned}$$

since s_{-q} is holomorphic by assumption. Therefore (3.2) shows that

$$(\nabla^{\theta})^* \nabla^{\theta} s = 2\Delta_{\bar{\partial}}^{\theta} s + i * F_{\nabla^{\theta}} s = ((2q + 1)\theta - q(q + 1)(2g - 2)) s.$$

The statement of (3.9) on the closure of this space of sections in $H^2(Z, \mathcal{E})$ follows by taking sequences of such sections and noting that the Laplacian is continuous as an operator $H^2(Z, \mathcal{E}) \rightarrow L^2(Z, \mathcal{E})$. \square

3.2. Spectral gaps around the low-lying eigenvalues. In this section we show that the eigensections of Proposition 3.6 are the only eigensections with eigenvalue bounded above by $\mu_m = (2m + 1)\theta - m(m + 1)(2g - 2)$. As a consequence, we see that in the interval $[0, \mu_m)$, the spectrum is discrete and consists of eigenvalues μ_0, \dots, μ_{m-1} .

Proposition 3.7. *In the interval $[0, \mu_m)$, the spectrum of the Laplacian $(\nabla^\theta)^* \nabla^\theta$ takes the discrete values $\mu_q = (2q + 1)\theta - q(q + 1)(2g - 2)$ for each $q = 0, \dots, m - 1$, and the space of eigensections is*

$$E_{\mu_q} = \overline{\partial^{\nabla^{\theta, -1}} \circ \dots \circ \partial^{\nabla^{\theta, -q}} (\ker \bar{\partial}^{\nabla^{\theta, -q}})} \subset \Omega^0(\tilde{\mathcal{E}}).$$

This defines an isomorphism of A_θ -modules $E_{\mu_q} \cong H^0(\tilde{\mathcal{E}} \otimes K^{-q})$.

Proof. From (3.2) it follows directly that $s \in \ker \bar{\partial}^{\nabla^\theta}$ implies that $(\nabla^\theta)^* \nabla^\theta s = \theta s = \mu_0 s$.

We have $\ker \bar{\partial}^{\nabla^\theta} = \ker(\partial^{\nabla^{\theta, -1}})^* = \text{coker } \partial^{\nabla^{\theta, -1}}$, and so there is a direct sum decomposition

$$\Omega^0(\tilde{\mathcal{E}}) \cong \ker \bar{\partial}^{\nabla^\theta} \oplus \overline{\text{im } \partial^{\nabla^{\theta, -1}}}.$$

First consider the case where $s \in \text{im } \partial^{\nabla^{\theta, -1}}$ (we will generalise this to $s \in \overline{\text{im } \partial^{\nabla^{\theta, -1}}}$ below). Write $s = \partial^{\nabla^{\theta, -1}} t$. We can make this choice unique by choosing $t \in (\ker \partial^{\nabla^{\theta, -1}})^\perp$, although this is not necessary in the following proof. Equation (3.8) shows that

$$\Delta_{\bar{\partial}}^\theta s = \Delta_{\bar{\partial}}^\theta \circ \partial^{\nabla^{\theta, -1}} t = i \partial^{\nabla^{\theta, -1}} * F_{\nabla^{\theta, -1}} t + \partial^{\nabla^{\theta, -1}} \circ \Delta_{\bar{\partial}}^{\theta, -1} t \in \text{im } \partial^{\nabla^{\theta, -1}}.$$

Since the curvature $i * F_{\nabla^{\theta, -1}}$ is constant, then $i \partial^{\nabla^{\theta, -1}} * F_{\nabla^{\theta, -1}} t = (i * F_{\nabla^{\theta, -1}}) \partial^{\nabla^{\theta, -1}} t = (i * F_{\nabla^{\theta, -1}}) s$, and so

$$(3.12) \quad (\nabla^\theta)^* \nabla^\theta s = 2\Delta_{\bar{\partial}}^\theta s + i * F_{\nabla^\theta} s = (2i * F_{\nabla^{\theta, -1}} + i * F_{\nabla^\theta}) s + 2\partial^{\nabla^{\theta, -1}} \circ \Delta_{\bar{\partial}}^{\theta, -1} t.$$

If $t \in \ker \bar{\partial}^{\nabla^{\theta, -1}}$, then the remaining term vanishes and we obtain

$$(3.13) \quad (\nabla^\theta)^* \nabla^\theta s = 2\Delta_{\bar{\partial}}^\theta s + i * F_{\nabla^\theta} s = (2i * F_{\nabla^{\theta, -1}} + i * F_{\nabla^\theta}) s,$$

which gives us the bound that we want, since $i * F_{\nabla^{\theta, -1}} = \theta - (2g - 2)$ is positive.

If $t \in (\ker \bar{\partial}^{\nabla^{\theta, -1}})^\perp$, then we want to show that the final term in (3.12) is non-negative. Take the L^2 inner product of both sides of (3.12) with $s = \partial^{\nabla^{\theta, -1}} t$ to obtain

$$(3.14) \quad \langle (\nabla^\theta)^* \nabla^\theta s, s \rangle = \langle (2i * F_{\nabla^{\theta, -1}} + i * F_{\nabla^\theta}) s, s \rangle + 2 \langle \partial^{\nabla^{\theta, -1}} \circ \Delta_{\bar{\partial}}^{\theta, -1} t, \partial^{\nabla^{\theta, -1}} t \rangle.$$

The final term is

$$\begin{aligned} \langle \partial^{\nabla^{\theta, -1}} \circ \Delta_{\bar{\partial}}^{\theta, -1} t, \partial^{\nabla^{\theta, -1}} t \rangle &= \langle \Delta_{\bar{\partial}}^{\theta, -1} \circ \Delta_{\bar{\partial}}^{\theta, -1} t, t \rangle \\ &= \langle (\Delta_{\bar{\partial}}^{\theta, -1} + i * F_{\nabla^{\theta, -1}}) \circ \Delta_{\bar{\partial}}^{\theta, -1} t, t \rangle \quad \text{from (3.1)} \\ &= \langle \Delta_{\bar{\partial}}^{\theta, -1} t, \Delta_{\bar{\partial}}^{\theta, -1} t \rangle + i * F_{\nabla^{\theta, -1}} \langle \Delta_{\bar{\partial}}^{\theta, -1} t, t \rangle. \end{aligned}$$

Since $i * F_{\nabla^{\theta, -1}} > 0$ and $\Delta_{\bar{\partial}}^{\theta, -1}$ is non-negative, then this final term from (3.14) is non-negative. Therefore (3.14) becomes

$$\begin{aligned} \langle (\nabla^\theta)^* \nabla^\theta s, s \rangle &\geq \langle (2i * F_{\nabla^{\theta, -1}} + i * F_{\nabla^\theta}) s, s \rangle \\ &= (2i * F_{\nabla^{\theta, -1}} + i * F_{\nabla^\theta}) \langle s, s \rangle \\ \Leftrightarrow \frac{\langle (\nabla^\theta)^* \nabla^\theta s, s \rangle}{\langle s, s \rangle} &\geq 2i * F_{\nabla^{\theta, -1}} + i * F_{\nabla^\theta}, \end{aligned}$$

and so there is a gap in the spectrum at $\theta = i * F_{\nabla^\theta}$, since $2i * F_{\nabla^{\theta, -1}} > 0$.

Now consider a general $s \in \overline{\text{im } \partial^{\nabla^{\theta,-1}}}$, and write $s = \lim_{n \rightarrow \infty} s_n$ for a sequence $\{s_n\} \subset \text{im } \partial^{\nabla^{\theta,-1}}$ such that $\|(\nabla^\theta)^* \nabla^\theta s_n\|$ is bounded and $\|s_n\|^2 = \|s\|^2$. As before, write $s_n = \partial^{\nabla^{\theta,-1}} t_n$ for each n and apply the same argument as above to show that

$$\frac{\langle (\nabla^\theta)^* \nabla^\theta s_n, s_n \rangle}{\langle s, s \rangle} = \frac{\langle (\nabla^\theta)^* \nabla^\theta s_n, s_n \rangle}{\langle s_n, s_n \rangle} \geq 2i * F_{\nabla^{\theta,-1}} + i * F_{\nabla^\theta}.$$

Therefore, since the operator $(\nabla^\theta)^* \nabla^\theta$ is continuous and $\|(\nabla^\theta)^* \nabla^\theta s_n\|$ is bounded, then

$$\frac{\langle (\nabla^\theta)^* \nabla^\theta s, s \rangle}{\langle s, s \rangle} = \lim_{n \rightarrow \infty} \frac{\langle (\nabla^\theta)^* \nabla^\theta s_n, s_n \rangle}{\langle s_n, s_n \rangle} \geq 2i * F_{\nabla^{\theta,-1}} + i * F_{\nabla^\theta}.$$

In summary, if $s \in \overline{\text{im } \partial^{\nabla^{\theta,-1}}}$, then $\frac{\langle (\nabla^\theta)^* \nabla^\theta s, s \rangle}{\langle s, s \rangle} \geq 2i * F_{\nabla^{\theta,-1}} + i * F_{\nabla^\theta}$, and so the eigenvalues of $(\nabla^\theta)^* \nabla^\theta$ on $\overline{\text{im } \partial^{\nabla^{\theta,-1}}}$ are bounded below by $2i * F_{\nabla^{\theta,-1}} + i * F_{\nabla^\theta}$. Since $2i * F_{\nabla^{\theta,-1}} > 0$ then there is a spectral gap at the lowest eigenvalue $i * F_{\nabla^\theta}$.

Continuing inductively, suppose that $s \in \overline{\text{im } \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-q}}}$. Again, we can decompose $\Omega^0(\tilde{\mathcal{E}} \otimes K^{-q}) \cong \overline{\text{im } \partial^{\nabla^{\theta,-(q+1)}}} \oplus \overline{\text{coker } \partial^{\nabla^{\theta,-(q+1)}}}$. Given any $t \in \Omega^0(\tilde{\mathcal{E}} \otimes K^{-q})$, there exists a sequence $\{t_n\} \subset \text{im } \partial^{\nabla^{\theta,-(q+1)}} \oplus \text{coker } \partial^{\nabla^{\theta,-(q+1)}}$ such that $\lim_{n \rightarrow \infty} t_n = t$. Write $t_n = \partial^{\nabla^{\theta,-(q+1)}} u_n + v_n$ with $u_n \in \Omega^0(\tilde{\mathcal{E}} \otimes K^{-(q+1)})$ and $v_n \in \text{coker } \partial^{\nabla^{\theta,-(q+1)}} \cong \ker \bar{\partial}^{\nabla^{\theta,-q}}$.

The same argument as before shows that

$$\begin{aligned} \Delta_{\bar{\partial}}^\theta (\partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-q}} v_n) &= \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-q}} \circ \Delta_{\bar{\partial}}^{\theta,-q} v_n \\ &\quad + \left(\sum_{\ell=1}^q (\theta - \ell(2g-2)) \right) \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-q}} v_n \\ &= \left(q\theta - \frac{q(q+1)(2g-2)}{2} \right) \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-q}} v_n, \end{aligned}$$

and therefore $\partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-q}} v_n$ is an eigensection of $(\nabla^\theta)^* \nabla^\theta = 2\Delta_{\bar{\partial}}^\theta + i * F_{\nabla^\theta}$ with eigenvalue $2 \left(q\theta - \frac{q(q+1)(2g-2)}{2} \right) + \theta = (2q+1)\theta - q(q+1)(2g-2)$.

Now consider $s_n = \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n$. Again, the same argument as above shows that

$$(3.15) \quad \Delta_{\bar{\partial}}^\theta s_n = \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n + \sum_{\ell=1}^{q+1} (\theta - \ell(2g-2)) s_n.$$

It remains to show that the inner product of the first term with $s_n = \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n$ is non-negative. To see this, write

$$\begin{aligned}
& \left\langle \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n \right\rangle \\
&= \left\langle (\partial^{\nabla^{\theta,-1}})^* \partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n \right\rangle \\
&= \left\langle \Delta_{\bar{\partial}}^{\theta,-1} \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n \right\rangle \\
&= \left\langle (i * F_{\nabla^{\theta,-1}} + \Delta_{\bar{\partial}}^{\theta,-1}) \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n \right\rangle \\
&= (i * F_{\nabla^{\theta,-1}}) \left\langle \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n \right\rangle \\
&\quad + \left\langle \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \Delta_{\bar{\partial}}^{\theta,-1} \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n \right\rangle.
\end{aligned}$$

In the final expression above, the first term is a positive multiple of

$$\left\langle \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n \right\rangle.$$

In order to evaluate the second term, we can use Lemma 3.5 and the same process as in the proof of Proposition 3.6 to write

$$\begin{aligned}
\Delta_{\bar{\partial}}^{\theta,-1} \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n &= \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n \\
&\quad + C \cdot \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n,
\end{aligned}$$

where C is a positive constant. Therefore we obtain a positive multiple of

$$\left\langle \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n \right\rangle$$

plus the non-negative term $\left\langle \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n \right\rangle$.

Now repeat the same process on the term

$$\left\langle \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, \partial^{\nabla^{\theta,-2}} \circ \dots \circ \partial^{\nabla^{\theta,-(q+1)}} u_n \right\rangle.$$

Continuing inductively, at the end of the process we obtain non-negative terms plus a positive multiple of the non-negative term $\left\langle \Delta_{\bar{\partial}}^{\theta,-(q+1)} u_n, u_n \right\rangle$. Substituting this into (3.15), we see that

$$\left\langle \Delta_{\bar{\partial}}^{\theta} s_n, s_n \right\rangle \geq \sum_{\ell=1}^{q+1} (\theta - \ell(2g - 2)) \langle s_n, s_n \rangle.$$

By taking sequences as before, this inequality is preserved in the limit, and therefore it applies to sections in the closure. Since $\theta - m(2g - 2) > 0$ by assumption, then the final term in this sum is strictly positive if $q \leq m - 1$, and so there is a gap in the spectrum of $\Delta_{\bar{\partial}}^{\theta}$ at the eigenvalue $\sum_{\ell=1}^q (\theta - \ell(2g - 2))$. By (3.2) there is then a gap in the spectrum of $(\nabla^{\theta})^* \nabla^{\theta}$ at the eigenvalue μ_q when $q \leq m - 1$.

Finally, since $\partial^{\nabla^{\theta,-\ell}}$ commutes with T_{γ} for all ℓ by the results of Section 1.3 and it is injective for all $\ell \leq m$ by Remark 3.1, then this $\partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-q}}$ defines an isomorphism

of A_θ modules

$$H^0(\tilde{\mathcal{E}} \otimes K^{-q}) \cong E_{\mu_q} = \overline{\partial^{\nabla^{\theta,-1}} \circ \dots \circ \partial^{\nabla^{\theta,-q}} (\ker \bar{\partial}^{\nabla^{\theta,-q}})}.$$

□

4. AN INDEX THEOREM

In the previous notation, let ∇^θ be the unitary connection on the trivial line bundle on Z , whose curvature is equal to $i\tilde{B} = i\theta\tilde{\omega}$. Consider the Dolbeault operator $\bar{\partial}$ on X , and lift it to $\tilde{\bar{\partial}}$ on Z , which is invariant under the group of deck transformations \mathbb{Z}^{2g} for the abelian cover $Z \rightarrow X$. Let $\mathcal{E} \rightarrow X$ be a holomorphic vector bundle over X , and $\nabla^\mathcal{E}$ a $(0,1)$ connection on \mathcal{E} . Let $\tilde{\mathcal{E}}$ denote the lift of \mathcal{E} to Z , and $\tilde{\nabla}^\mathcal{E}$ the lift of $\nabla^\mathcal{E}$ to Z . Then as before, one checks that the operator $\tilde{\bar{\partial}} \otimes \tilde{\nabla}^\mathcal{E} \otimes \nabla^\theta$ is invariant under the projective action of \mathbb{Z}^{2g} as in the previous section. Here

$$(4.1) \quad \tilde{\bar{\partial}} \otimes \tilde{\nabla}^\mathcal{E} \otimes \nabla^\theta : \Omega_{(2)}^{0,0}(Z, \mathcal{E}) \longrightarrow \Omega_{(2)}^{0,1}(Z, \mathcal{E})$$

where $\Omega_{(2)}^{0,j}(Z, \mathcal{E})$, $j = 0, 1$ denotes the space of square integrable differential j -forms on Z with coefficients in \mathcal{E} .

Define

$$h^0(\tilde{\nabla}^\mathcal{E} \otimes \nabla^\theta) = \dim_\tau(\ker_{L^2}(\tilde{\bar{\partial}} \otimes \tilde{\nabla}^\mathcal{E} \otimes \nabla^\theta)).$$

By L^2 -Serre duality,

$$h^1(\tilde{\nabla}^\mathcal{E} \otimes \nabla^\theta) = h^0(\tilde{\nabla}^{\mathcal{E}^* \otimes K} \otimes \nabla^{-\theta}) = \dim_\tau(\ker_{L^2}(\tilde{\bar{\partial}} \otimes \tilde{\nabla}^{\mathcal{E}^* \otimes K} \otimes \nabla^{-\theta})).$$

where τ denotes the von Neumann trace or dimension. Elliptic regularity ensures that these numbers are finite.

The goal of this section is to prove the following result.

Theorem 4.1 (L^2 -Riemann-Roch for projective actions). *The L^2 index of the induced connection $\tilde{\bar{\partial}} \otimes \nabla^{K^{-j}} \otimes \nabla^\theta$ is*

$$(4.2) \quad \begin{aligned} \text{index}_{L^2}(\tilde{\bar{\partial}} \otimes \nabla^\mathcal{E} \otimes \nabla^\theta) &= h^0(\nabla^\mathcal{E} \otimes \nabla^\theta) - h^1(\nabla^\mathcal{E} \otimes \nabla^\theta) \\ &= \deg(\mathcal{E}) + rk(\mathcal{E})\theta + (1-g)rk(\mathcal{E}) \end{aligned}$$

In the case where $\mathcal{E} = K_X^{-q}$ is a power of the canonical bundle on X , then

(1) the L^2 index is

$$h^0(\nabla^{K^{-q}} \otimes \nabla^\theta) - h^1(\nabla^{K^{-q}} \otimes \nabla^\theta) = (2q+1)(1-g) + \theta.$$

(2) (vanishing) If $\theta > 2g - 2$, then $h^1(\nabla^\theta) = 0$, so that $h^0(\nabla^\theta) = 1 - g + \theta$.

Proof. Note that since Z is a Riemann surface, then (4.1) is a complex, and by the higher twisted index theorem in [18], the index is

$$\text{Index}_{A_\theta}(\tilde{\bar{\partial}} \otimes \tilde{\nabla}^\mathcal{E} \otimes \nabla^\theta) = [\ker(\tilde{\bar{\partial}} \otimes \tilde{\nabla}^\mathcal{E} \otimes \nabla^\theta) - \text{coker}(\tilde{\bar{\partial}} \otimes \tilde{\nabla}^\mathcal{E} \otimes \nabla^\theta)] \in K_0(\mathcal{A}_\theta).$$

As in the index for a family of elliptic operators, in general neither $\text{coker}(\tilde{\partial} \otimes \tilde{\nabla}^\varepsilon \otimes \nabla^\theta)$ nor $\text{ker}(\tilde{\partial} \otimes \tilde{\nabla}^\varepsilon \otimes \nabla^\theta)$ are finite projective modules, however under a vanishing condition $\text{coker}(\tilde{\partial} \otimes \tilde{\nabla}^\varepsilon \otimes \nabla^\theta) = 0$, then $\text{ker}(\tilde{\partial} \otimes \tilde{\nabla}^\varepsilon \otimes \nabla^\theta)$ is a finite projective module, which is therefore a holomorphic vector bundle in terminology of subsection 1.5.

Then by the twisted index theorem [21], one has

$$\begin{aligned} \tau \left(\text{Index}_{A_\theta}(\tilde{\partial} \otimes \tilde{\nabla}^\varepsilon \otimes \nabla^\theta) \right) &= \frac{1}{2\pi} \int_X \text{Todd}(\Omega_X) \text{tr} \exp(\Omega^\varepsilon) \exp(\theta\omega) \\ &= \frac{1}{2\pi} \int_X \left(1 + \frac{1}{2}\Omega_X\right) (rk(\mathcal{E}) + \text{tr}(\Omega^\varepsilon))(1 + \theta\omega) \\ &= \text{deg}(\mathcal{E}) + rk(\mathcal{E})\theta + (1 - g)rk(\mathcal{E}). \end{aligned}$$

On the other hand,

$$\tau \left(\text{Index}_{A_\theta}(\tilde{\partial} \otimes \tilde{\nabla}^\varepsilon \otimes \nabla^\theta) \right) = h^0(\tilde{\nabla}^\varepsilon \otimes \nabla^\theta) - h^1(\tilde{\nabla}^\varepsilon \otimes \nabla^\theta),$$

which completes the proof of (4.2).

Now let $\mathcal{E} = K_X^{-q}$ be the holomorphic line bundle. We conclude from (4.2) that

$$(4.3) \quad h^0(\tilde{\nabla}^{K^{-q}} \otimes \nabla^\theta) = (-q \text{deg}(K_X) + \theta) + 1 - g = (2q + 1)(1 - g) + \theta > 0.$$

If $q \geq 0$ is such that $(q + 1)(2g - 2) - \theta < 0$, then (3.1) implies that $h^0(\tilde{\nabla}^{K^{q+1}} \otimes \nabla^{-\theta}) = 0$. Using L^2 Serre duality we conclude that

$$(4.4) \quad h^1(\tilde{\nabla}^{K^{-q}} \otimes \nabla^\theta) = h^0(\tilde{\nabla}^{K^{q+1}} \otimes \nabla^{-\theta}) = 0.$$

□

Theorem 3.2 shows that $\text{ker}_{A_\theta}(\tilde{\partial} \otimes \tilde{\nabla}^\varepsilon \otimes \nabla^\theta) = E_{\mu_q}$, the μ_q -th eigenspace of the magnetic Laplacian. Using Theorem 4.1 and the terminology in subsection 1.5, we have the following result on the dimension of these eigenspaces, which proves part (3) of Theorem 2.1.

Lemma 4.2. *Let E_{μ_q} be the μ_q -th eigenspace of the magnetic Laplacian. Then*

- (1) E_{μ_q} is a holomorphic vector bundle over $\mathcal{A}_\theta \otimes \mathcal{K}(L^2(\mathcal{F}))$, and
- (2) The von Neumann dimension is $\dim_\tau(E_{\mu_q}) = (2q + 1)(1 - g) + \theta > 0$.

5. CHERN NUMBER OF E_{μ_q} VIA A HIGHER INDEX THEOREM

Our goal is to calculate the Chern number of the spectral subspace E_{μ_q} as a finite projective A_θ -module, which will complete the proof of the final part of Theorem 2.1. We do this by using the higher twisted index theorem [18, Thm. 2.2]. The argument goes as follows. Let c be the group cocycle on \mathbb{Z}^{2g} corresponding to the symplectic 2-form, and let τ_c be the corresponding continuous 2-cocycle from (1.6). Applying the higher twisted index theorem and simplifying, we see that

$$\tau_c(\text{index}_{A_\theta}(\tilde{\partial} \otimes \tilde{\nabla}^{K^{-j}} \otimes \nabla^\theta)) = \int_X \phi(c).$$

By the vanishing theorem (4.4), we deduce that if $\theta > (j+1)(2g-2)$ then $\ker_{A_\theta}(\tilde{\partial} \otimes \tilde{\nabla}^{K^{-j}} \otimes \nabla^\theta)$ is a finite projective A_θ module, and

$$\tau_c(\ker_{A_\theta}(\tilde{\partial} \otimes \tilde{\nabla}^{K^{-j}} \otimes \nabla^\theta)) = \int_X \phi(c).$$

Now $\phi(c)$ is easily seen to be the Bergman volume form on X , and $\int_X \phi(c) = 2g$ if X is a genus g Riemann surface. Finally, we will prove in Theorem 3.2 that

$$\ker_{A_\theta}(\tilde{\partial} \otimes \tilde{\nabla}^{K^{-j}} \otimes \nabla^\theta) \cong E_{\mu_q}$$

as A_θ modules, therefore we conclude that if X is a genus g Riemann surface, then the Chern number of E_{μ_q} is

$$(5.1) \quad \tau_c(E_{\mu_q}) = \int_\Sigma \phi(c) = 2g,$$

which proves (2.1).

We can extend the above calculations to the case of a “good” orbifold given by a quotient $X = X'/\Gamma$ as follows. Let g' be the genus of X' and let $p : X' \rightarrow X$ be the associated ramified cover. Since $\phi(c)$ pulls back to the Bergman volume form on X' , then

$$\int_X \phi(c) = \frac{1}{|\Gamma|} \int_{X'} p^* \phi(c) = \frac{2g'}{|\Gamma|}.$$

Let $n_y = |\Gamma_y|$ be the order of the isotropy group at each point $y \in X'$, let $R = \{y \in X' : n_y > 1\}$ be the ramification divisor, and let $n = |R|$ be the total number of ramification points in X' . The Riemann-Hurwitz theorem shows that

$$\begin{aligned} 2g' - 2 &= |\Gamma|(2g - 2) + \sum_{y \in X'} (n_y - 1) \\ \Leftrightarrow \frac{2g'}{|\Gamma|} &= 2g - 2 + \frac{2}{|\Gamma|} + \frac{1}{|\Gamma|} \sum_{y \in X'} (n_y - 1) \\ &= 2g - 2 + \frac{1}{|\Gamma|} \sum_{y \in R} n_y + \frac{2 - n}{|\Gamma|}. \end{aligned}$$

The term $\frac{1}{|\Gamma|} \sum_{y \in R} n_y$ can be further simplified by noting that for each $y \in R$, the number of points in the orbit $\Gamma \cdot y$ is equal to $\frac{|\Gamma|}{n_y}$. Therefore $\frac{1}{|\Gamma|} \sum_{y \in R} n_y$ is equal to the number of Γ -orbits in R , which we denote by $\#(R/\Gamma)$. In conclusion, we have proved that

$$\int_X \phi(c) = 2g - 2 + \#(R/\Gamma) + \frac{2 - n}{|\Gamma|}.$$

By Theorem 3.2, we know that there is an isomorphism of A_θ modules $\ker_{A_\theta}(\tilde{\partial} \otimes \tilde{\nabla}^\varepsilon \otimes \nabla^\theta) \cong E_{\mu_q}$, the μ_q -th eigenspace of the magnetic Laplacian. All of the above is summarised in the following lemma, which proves (2.2) and thus concludes the proof of Theorem 2.1.

Lemma 5.1. *The Chern class of the μ_q -th eigenspace of the magnetic Laplacian is*

$$(5.2) \quad \tau_c(E_{\mu_q}) = 2g - 2 + \#(R/\Gamma) + \frac{2-n}{|\Gamma|}.$$

REFERENCES

- [1] L. Alvarez-Gaumé, G. Moore, and C. Vafa. Theta functions, modular invariance, and strings. *Comm. Math. Phys.*, 106(1):1–40, 1986.
- [2] J. E. Avron, R. Seiler, and L. G. Yaffe. Adiabatic theorems and applications to the quantum Hall effect. *Comm. Math. Phys.*, 110(1):33–49, 1987.
- [3] J. E. Avron, R. Seiler, and P. G. Zograf. Adiabatic quantum transport: Quantization and fluctuations. *Phys. Rev. Lett.*, 73:3255–3257, Dec 1994.
- [4] J. Bellissard, A. van Elst, and H. Schulz-Baldes. The noncommutative geometry of the quantum Hall effect. *J. Math. Phys.* volume 35, pages 5373–5451. 1994.
- [5] M. T. Benameur and V. Mathai. Gap-labelling conjecture with nonzero magnetic field. *Adv. Math.*, 325:116–164, 2018.
- [6] J. Brüning and T. Sunada. On the spectrum of gauge-periodic elliptic operators. Number 210, pages 6, 65–74. 1992. *Méthodes semi-classiques*, Vol. 2 (Nantes, 1991).
- [7] A. L. Carey, K. C. Hannabuss, and V. Mathai. Quantum Hall effect on the hyperbolic plane in the presence of disorder. *Lett. Math. Phys.*, 47(3):215–236, 1999.
- [8] A. L. Carey, K. C. Hannabuss, V. Mathai, and P. McCann. Quantum Hall effect on the hyperbolic plane. *Comm. Math. Phys.*, 190(3):629–673, 1998.
- [9] A. L. Carey, K. C. Hannabuss, and V. Mathai. Quantum Hall effect and noncommutative geometry. *J. Geom. Symmetry Phys.*, 6:16–37, 2006.
- [10] A. Comtet and P. J. Houston. Effective action on the hyperbolic plane in a constant external field. *J. Math. Phys.*, 26(1):185–191, 1985.
- [11] G. A. Elliott. On the K -theory of the C^* -algebra generated by a projective representation of a torsion-free discrete abelian group. In *Operator algebras and group representations, Vol. I (Neptun, 1980)*, volume 17 of *Monogr. Stud. Math.*, pages 157–184. Pitman, Boston, MA, 1984.
- [12] O. Forster. *Lectures on Riemann surfaces*, volume 81 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1981. Translated from the German by B. Gilligan.
- [13] H. B. Griffiths. The fundamental group of a surface, and a theorem of Schreier. *Acta Math.*, 110:1–17, 1963.
- [14] D. Hilbert. Ueber Flächen von constanter Gausscher Krümmung. *Trans. Amer. Math. Soc.*, 2(1):87–99, 1901.
- [15] S. Klevtsov, X. Ma, G. Marinescu, and P. Wiegmann. Quantum Hall effect and Quillen metric. *Comm. Math. Phys.*, 349(3):819–855, 2017.
- [16] A. J. Kollár, M. Fitzpatrick, and A. A. Houck. Hyperbolic lattices in circuit quantum electrodynamics. *Nature*, 571(7763):45–50, 2019.
- [17] M. Marcolli and V. Mathai. Twisted index theory on good orbifolds. I. Noncommutative Bloch theory. *Commun. Contemp. Math.*, 1(4):553–587, 1999.
- [18] M. Marcolli and V. Mathai. Twisted index theory on good orbifolds. II. Fractional quantum numbers. *Comm. Math. Phys.*, 217(1):55–87, 2001.
- [19] M. Marcolli and V. Mathai. Towards the fractional quantum Hall effect: a noncommutative geometry perspective. In *Noncommutative geometry and number theory*, *Aspects Math.*, E37, pages 235–261. Friedr. Vieweg, Wiesbaden, 2006.
- [20] M. Marcolli and K. Seipp. Twisted index theory on orbifold symmetric products and the fractional quantum Hall effect. *Adv. Theor. Math. Phys.*, 21(2):451–501, 2017.
- [21] V. Mathai. K -theory of twisted group C^* -algebras and positive scalar curvature. In *Tel Aviv Topology Conference: Rothenberg Festschrift (1998)*, volume 231 of *Contemp. Math.*, pages 203–225. Amer. Math. Soc., Providence, RI, 1999.
- [22] V. Mathai and D. Quillen. Superconnections, Thom classes, and equivariant differential forms. *Topology*, 25(1):85–110, 1986.

- 1
2
3 [23] V. Mathai and J. Rosenberg. The Riemann-Roch theorem on higher dimensional complex noncommu-
4 tative tori. *J. Geom. Phys.*, 147:103534, 9, 2020.
- 5 [24] V. Mathai and G. C. Thiang. Topological phases on the hyperbolic plane: fractional bulk-boundary
6 correspondence. *Adv. Theor. Math. Phys.*, 23(3):803–840, 2019.
- 7 [25] V. Mathai and G. Wilkin. Fractional quantum numbers via complex orbifolds. *Lett. Math. Phys.*,
8 109(11):2473–2484, 2019.
- 9 [26] E. Prodan and H. Schulz-Baldes. *Bulk and boundary invariants for complex topological insulators*. Math-
10 ematical Physics Studies. Springer, [Cham], 2016. From K -theory to physics.
- 11 [27] A. Putman. Cutting and pasting in the Torelli group. *Geom. Topol.*, 11:829–865, 2007.
- 12 [28] D. Quillen. Determinants of Cauchy-Riemann operators on Riemann surfaces. *Funktsional. Anal. i*
13 *Prilozhen.*, 19(1):37–41, 96, 1985.
- 14 [29] C. Tejero Prieto. Fourier-Mukai transform and adiabatic curvature of spectral bundles for Landau
15 Hamiltonians on Riemann surfaces. *Comm. Math. Phys.*, 265(2):373–396, 2006.
- 16 [30] C. Tejero Prieto. Holomorphic spectral geometry of magnetic Schrödinger operators on Riemann sur-
17 faces. *Differential Geom. Appl.*, 24(3):288–310, 2006.
- 18 [31] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs. Quantized hall conductance in a
19 two-dimensional periodic potential. *Phys. Rev. Lett.*, 49:405–408, Aug 1982.
- 20

21 SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF ADELAIDE, ADELAIDE SA 5005 AUSTRALIA.
22 *E-mail address:* mathai.varghese@adelaide.edu.au

23
24 DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YO10 5DD, UK
25 *E-mail address:* graeme.wilkin@york.ac.uk

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60