

09 PM
M4608



Bacteriophage SfII Mediated Serotype Conversion in *Shigella flexneri*

by

Maria Mavris B.Sc.(Hons)(Adel)




Microbial Pathogenesis Unit
Department of Microbiology and Immunology
University of Adelaide
Adelaide 5005
Australia

A Thesis Submitted for the Degree of Doctor of Philosophy
May, 1998

Abstract

Shigella flexneri is the causative agent of bacillary dysentery and is responsible for approximately 10^5 deaths per year. Organisms that belong to this species can be classified according to the O-antigenic component of their lipopolysaccharide (LPS). Determination of the serotype is based on agglutination of the organism with polyclonal antisera. *S. flexneri* exhibits at least 12 different serotypes, all of which are variations of the basic tetrasaccharide repeat:



I II III

Variations resulting in different epitopes are due to the addition of either glucosyl or O-acetyl residues at specific positions along the repeat unit. Immunity to *S. flexneri* strains is serotype specific and therefore protection against each serotype is required.

The modifications have been found to be mediated by temperate bacteriophages. To date, two such phages have been identified and characterised; Sf6, responsible for the O-acetylation of rhamnose III and SfX which encodes genes which add a glucosyl residue to rhamnose I.

S. flexneri strains of serotype 2 have been associated with a highly virulent phenotype and are more prevalent in nature. Bacteriophage SfII is responsible for the glucosylation of rhamnose III of the repeat unit and has been isolated from a strain of serotype 2b. Phage SfII possesses an hexagonal shaped head, a tail with a contractile sheath and tail fibres and has a genome of 43.5 kb in size.

A 4 kb *Bam*HI fragment was isolated which was found to mediate serotype conversion. This fragment was found to contain 4 open reading frames, however, only two were required for serotype conversion. These genes were named *bgt*, which encodes a putative bactoprenol glucosyl transferase, and *gtrII* encoding the putative type II antigen determining glucosyl

transferase. These genes are adjacent to the integrase and excisionase genes and the attachment site (*attP*) which are highly homologous to those of *Salmonella* bacteriophage P22. In addition, a gene named ORF2 was identified of unknown function, which did not appear to be required for serotype conversion.

In vitro T7 polymerase /promoter analysis identified a protein of 34 kDa in size corresponding to Bgt. Subsequent subcellular fractionation localised this protein to the cytoplasmic and membrane fractions. Topology studies determined that the carboxy and amino terminal ends were cytoplasmically located. The active site was located to the cytoplasmic domain which is consistent with the role of Bgt. A protein was not observed for GtrII.

The ability of SfII phage and plasmid copies of serotype converting genes to modify LPS with only a single repeat unit was assessed. O-antigen polymerase mutants (*rfc*) of serotype X and Y, were transformed with various serotype converting plasmids and characterised by colony immunoblotting for reactivity to a variety of type and group antisera. In *S. flexneri*, it appears that a single O-antigen repeat unit is sufficient to be modified.

TABLE OF CONTENTS

1. INTRODUCTION	1
1.1 INTRODUCTION	1
1.2 DISCOVERY OF SHIGELLA SPECIES	1
1.2.1 Early serotyping of <i>Shigella flexneri</i> species	2
1.2.2 Work of J.S. Boyd	3
1.3 SHIGELLA SPECIES	3
1.3.1 Epidemiology	4
1.4 LIFESTYLE OF <i>S. FLEXNERII</i>	5
1.5 GENETICS OF SHIGELLA VIRULENCE.....	5
1.5.1 <i>Chromosomal loci</i>	6
1.5.1.1 <i>virR</i> , a central regulatory locus	6
1.5.1.2 Virulence associated chromosomal (<i>vac</i>) genes	7
1.5.1.3 Outer membrane protein C (OmpC)	8
1.6 PLASMID MEDIATED FUNCTIONS	8
1.6.1 Adhesion of bacteria to cell surface	9
1.6.2 Entry by induced phagocytosis.....	9
1.6.3 Escape from the phagocytic vacuole	10
1.6.4 Intracellular multiplication and spread.....	10
1.6.5 Lysis of double membrane surrounding protrusion.....	11
1.6.6 Biology of IcsA	11
1.7 LIPOPOLYSACCHARIDE (LPS)	13
1.7.1 Biosynthesis of LPS	14
1.7.2 Lipid A- genetics and biosynthesis.....	15
1.7.3 Core region.....	16
1.7.3.1 Inner core: 3-deoxy-D-mannoctulosonic acid (KDO).....	17
1.7.3.2 Outer core.....	17
1.7.4 O-antigen	18
1.7.4.1 Genetics of O antigen.....	19
1.7.4.2 O-polysaccharide biosynthesis.....	19
1.7.4.2.1 Enterobacterial Common Antigen (ECA)	21
1.7.4.3 Other genes required for O antigen synthesis	21
1.7.4.4 RfbX	22
1.7.4.5 RfaL, O-antigen ligase	22
1.7.4.6 Rfc, O-antigen polymerase.....	22
1.7.4.7 Rol, regulator of O antigen chain length.....	23
1.8 LIPOPOLYSACCHARIDE AND THE DISTRIBUTION OF ICSA	24
1.9 CHEMICAL UNDERSTANDING OF O-ANTIGEN OF <i>S. FLEXNERI</i>	26
1.10 VARIATION IN O ANTIGEN.....	27
1.10.1 <i>Vibrio cholerae</i> and serotype conversion	27
1.10.1.1 <i>V. cholerae</i> O139	28
1.10.2 <i>Salmonella enterica</i> serovar Typhimurium and serotypic variation	29
1.11 BACTERIOPHAGES WHICH HAVE THEIR RECEPTOR IN THE O-ANTIGEN	30
1.11.1 Lysogenisation of group E <i>Salmonellae</i>	31
1.11.2 Serotype converting bacteriophages of <i>S. flexneri</i>	32
1.11.2.1 Identification of <i>S. flexneri</i> bacteriophages.....	33

1.11.2.2 Identification of bacteriophages fII , f7,8 and fV.....	33
1.11.2.3 Bacteriophage Sf6.....	34
1.11.3 Molecular basis for serotype conversion.....	35
1.12 AIMS OF THIS STUDY:	35
2. MATERIALS AND METHODS	36
2.1 BACTERIAL STRAINS, BACTERIOPHAGES AND PLASMIDS	36
2.2 MAINTENANCE OF BACTERIAL AND BACTERIOPHAGE STRAINS.....	36
2.3 GROWTH MEDIA.....	37
2.4 CHEMICALS AND REAGENTS.....	38
2.5 ENZYMES AND IMMUNOCONJUGATES	39
2.6 TRANSFORMATION PROCEDURE	39
2.6.1 Super competent cells.....	40
2.7 ELECTROPORATION OF <i>S. FLEXNERI</i> AND <i>E. COLI</i>	40
2.8 BACTERIAL CONJUGATION	41
2.9 PLASMID DNA EXTRACTION PROCEDURES	41
2.10 PREPARATION OF <i>S. FLEXNERI</i> OR <i>E. COLI</i> GENOMIC DNA.....	44
2.10.1 Quick method for genomic DNA extraction	44
2.11 ANALYSIS AND MANIPULATION OF DNA	45
2.11.1 DNA quantitation	45
2.11.2 Restriction endonuclease digestion of DNA	45
2.11.3 Analytical and preparative separation of restriction fragments.....	45
2.11.4 Calculation of restriction fragment size	46
2.12 DNA CLONING PROCEDURES	47
2.12.1 Dephsophorylation of DNA using alkaline phosphatase.....	47
2.12.2 Ligation of DNA	47
2.12.3 <i>In vitro</i> cloning	47
2.12.4 Cosmid Cloning.....	48
2.13 SEQUENCING METHODS.....	48
2.13.1 Sequencing using dye labelled primers.....	48
2.13.2 Sequencing with dye-labelled terminators	49
2.13.3 Analysis of DNA sequences.....	49
2.14 POLYMERASE CHAIN REACTION PROTOCOL (PCR)	50
2.14.1 Synthesis of oligodeoxynucleotides	50
2.15 SOUTHERN TECHNIQUES	51
2.15.1 Preparation of DIG-labelled DNA probes.....	51
2.15.1.1 DIG labelled PCR	51
2.15.2 Southern transfer and hybridisation.....	52
2.16 SDS-POLYACRYLAMIDE GEL ELECTROPHORESIS (SDS-PAGE) AND WESTERN TRANSFER.	52
2.16.1 SDS-PAGE.....	52
2.16.2 Western transfer and detection	53
2.17 LIPOPOLYSACCHARIDE (LPS) PREPARATION.....	54
2.17.1 LPS-specific silver staining.....	54
2.17.2 Colony immunoblotting of LPS	54
2.17.3 Bacterial agglutination assays	55
2.17.4 O antigen hydrolysis assay	55
2.17.4.1 Preparation of formaldehyde (formalin)-fixed cells.....	55
2.17.4.2 O antigen hydrolysis assay.....	55
2.18 T7 POLYMERASE / PROMOTER EXPRESSION SYSTEM AND L-[³⁵ S]-METHIONINE LABELLING	56

2.18.1 Cell Fractionation	57
2.18.2 Autoradiography.....	57
2.19 ALKALINE PHOSPHATASE AND B-GALACTOSIDE ASSAYS	58
2.19.1 Alkaline phosphatase assays.....	58
2.19.2 β -galactosidase assays.....	58
2.20 BACTERIOPHAGE TECHNIQUES.....	59
2.20.1 Construction of lysogens	59
2.20.2 Phage assays.....	59
2.20.3 Preparation of phage stocks.....	60
2.20.3.1 Low titre phage stocks	60
2.20.3.2 High titre phage stocks by liquid infection	60
2.20.3.3 CsCl block density gradient for preparation of high titre phage stocks.....	61
2.20.3.4 Phenol extraction of bacteriophage DNA	61
2.20.3.5 Bacteriophage sensitivity tests	62
2.21 ELECTRON MICROSCOPY (EM)	62
3. ISOLATION AND CHARACTERISATION OF <i>S. FLEXNERI</i> SEROTYPE CONverting BACTERIOPHAGE SFII.	63
3.1 INTRODUCTION	63
3.2 RESULTS	64
3.2.1 Attempts to isolate bacteriophage SfII.....	64
3.2.2 Cosmid cloning of serotype conversion genes from strain PE655.....	64
3.2.3 Isolation of bacteriophage SfII from NCTC4.....	65
3.2.4 Purification and characterisation of bacteriophage SfII.....	65
3.2.4.1 Proteins of bacteriophage SfII.....	66
3.2.4.2 O-antigen hydrolysis assays.....	66
3.2.4.3 SfII lysogenisation of <i>rfa</i> mutant RMM109 of serotype Y	67
3.2.5 Characterisation of lysogens of bacteriophage SfII.....	67
3.2.6 Isolation and cloning of <i>Pst</i> I fragments of SfII genome.....	68
3.2.7 Characterisation of <i>Pst</i> I fragments of SfII genome.....	69
3.2.8 Identification of a region of bacteriophage SfII with serotype-converting ability.....	70
3.3 SUMMARY	70
4. SEQUENCE OF PRMM264 AND IDENTIFICATION OF THE ATTACHMENT REGION OF BACTERIOPHAGE SFII.....	71
4.1 INTRODUCTION	71
4.2 RESULTS	72
4.2.1 DNA sequencing of the <i>Bam</i> HI- <i>Bam</i> HI insert in pRMM264	72
4.2.2 Open reading frames in 4 kb <i>Bam</i> HI fragment of pRMM264	72
4.2.3 Open reading frames in 5.2 kb sequence.....	73
4.2.4 Integrase (<i>int</i>) and Excisionase (<i>xis</i>).....	73
4.2.5 The attachment site, <i>attP</i>	73
4.3 SUMMARY	75
5. LOCALISATION OF THE SEROTYPE CONVERTING GENES OF BACTERIOPHAGE SFII.....	76
5.1 INTRODUCTION	76
5.2 RESULTS	77
5.2.1 ORF2	77

5.2.2 ORF3 bactoprenol glucosyl transferase (<i>bgt</i>).....	77
5.2.3 ORF4 - glucosyl transferase <i>gtrII</i>	78
5.3 COMPLEMENTATION ANALYSES	79
5.3.1 Serotype conversion in <i>S. flexneri</i> with plasmid constructs	79
5.3.2 Serotype conversion with <i>gtrX</i> and <i>bgt</i>	79
5.3.3 Serotype conversion using <i>E. coli</i> K-12 DH5 α ORF3 homologue, <i>o306</i>	80
5.4. SEROTYPE CONVERSION IN <i>S. FLEXNERI</i> RFC MUTANTS.....	80
5.4.1 <i>rfc</i> mutants of <i>S. flexneri</i> strains of serotype Y and X.....	81
5.4.2 Serotype conversion of spontaneous <i>rfc</i> mutants	81
5.5 DISTRIBUTION OF BGT AND GTRII IN <i>S. FLEXNERI</i>	82
5.6 COMPARISON OF GENOME ORGANISATIONS IN PHAGE P22 AND <i>E. COLI</i> K-12	83
5.7 SUMMARY	83
6. CHARACTERISATION OF <i>S. FLEXNERI</i> TYPE II NEGATIVE MUTANTS AND BGT PROTEIN ANALYSES.....	85
6.1 INTRODUCTION.....	85
6.2 RESULTS	86
6.2.1 Isolation of type II antigen negative mutants of <i>S. flexneri</i>	86
6.2.2 Characterisation of type II antigen negative derivatives of RMM254	86
6.2.2.1 Inverse PCR of <i>TnphoA</i> insertion in RMA901	88
6.2.3 Complementation studies	89
6.2.3.1 LPS and western immunoblot analyses of complemented mutants RMA901 and RMA903	90
6.3 DETECTION OF BGT AND GTRII GENE PRODUCTS	91
6.3.1 Cellular location of Bgt in <i>E. coli</i>	92
6.3.2 Studies on the topology of Bgt using <i>phoA</i> and <i>lacZ</i> fusions.....	92
6.3.2.1 Tm-predict model of Bgt topology	93
6.3.2.2 Western immunoblotting of Bgt::PhoA and Bgt::LacZ fusions	94
6.4 SUMMARY AND CONCLUSIONS	95
7. DISCUSSION	96
7.1. INTRODUCTION	96
7.2 BACTERIOPHAGE SFII AND OTHER SEROTYPE DETERMINING BACTERIOPHAGES.....	97
7.3 LYSOGENS OF BACTERIOPHAGE SFII.....	99
7.4 GENOME MAPPING OF BACTERIOPHAGE SFII.....	100
7.5 BACTERIOPHAGE SFII ATTACHMENT SITE.....	101
7.6 A TWO STEP PATHWAY FOR SEROTYPE CONVERSION.....	102
7.7 DPM1- DOLICHOL PHOSPHATE MANNOSYL SYNTHASE.....	103
7.8 COMPLEMENTATION ANALYSES	104
7.9 COMPARISON OF ORGANISATION OF GENOMES	106
7.10 BGT AND GTRII PROTEINS	106
7.11 ADDITION OF MODIFICATION.....	107
7.12 CONCLUSIONS.....	108
8. BIBLIOGRAPHY	110