

References

Adachi W., Okubo K., Kinoshita S. (2000). Human uroplakin Ib in ocular surface epithelium. *Invest Ophthalmol Vis Sci* **41**: 2900-2905

Adams M.D., Celiker S.E., Holt R.A., Evans C.A., Gocayne J.D., Amanatides P.G., Scherer S.E., Li P.W., Hoskins R.A., Galle R.F., George R.A., Lewis S.E., Richards S., Ashburner M., Henderson S.N., Sutton G.G., Wortman J.R., Yandell M.D., Zhang Q., Chen L.X., Brandon R.C., Rogers Y.H., Blazej R.G., Champe M., Pfeiffer B.D., Wan K.H., Doyle C., Baxter E.G., Helt G., Nelson C.R., Gabor G.L., Abril J.F., Agbayani A., An H.J., Andrews-Pfannkoch C., Baldwin D., Ballew R.M., Basu A., Baxendale J., Bayraktaroglu L., Beasley E.M., Beeson K.Y., Benos P.V., Berman B.P., Bhandari D., Bolshakov S., Borkova D., Botchan M.R., Bouck J., Brokstein P., Brottier P., Burtis K.C., Busam D.A., Butler H., Cadieu E., Center A., Chandra I., Cherry J.M., Cawley S., Dahlke C., Davenport L.B., Davies P., de Pablos B., Delcher A., Deng Z., Mays A.D., Dew I., Dietz S.M., Dodson K., Doucette L.E., Downes M., Dugan-Rocha S., Dunkov B.C., Dunn P., Durbin K.J., Evangelista C.C., Ferraz C., Ferriera S., Fleischmann W., Fosler C., Gabrielian A.E., Garg N.S., Gelbart W.M., Glasser K., Glodek A., Gong F., Gorrell J.H., Gu Z., Guan P., Harris M., Harris N.L., Harvey D., Heiman T.J., Hernandez J.R., Houck J., Hostin D., Houston K.A., Howland T.J., Wei M.H., Ibegwam C., Jalali M., Kalush F., Karpen G.H., Ke Z., Kennison J.A., Ketchum K.A., Kimmel B.E., Kodira C.D., Kraft C., Kravitz S., Kulp D., Lai Z., Lasko P., Lei Y., Levitsky A.A., Li J., Li Z., Liang Y., Lin X., Liu X., Mattei B., McIntosh T.C., McLeod M.P., McPherson D., Merkulov G., Milshina N.V., Mobarry C., Morris J., Moshrefi A., Mount S.M., Moy M., Murphy B., Murphy L., Muzny D.M., Nelson D.L., Nelson D.R., Nelson K.A., Nixon K., Nusskern D.R., Pacleb J.M., Palazzolo M., Pittman G.S., Pan S., Pollard J., Puri V., Reese M.G., Reinert K., Remington K., Saunders R.D., Scheeler F., Shen H., Shue B.C., Siden-Kiamos I., Simpson M., Skupski M.P., Smith T., Spier E., Spradling A.C., Stapleton M., Strong R., Sun E., Svirskas R., Tector C., Turner R., Venter E., Wang A.H., Wang X., Wang Z.Y., Wassarman D.A., Weinstock G.M., Weissenbach J., Williams S.M., Woodage T., Worley K.C., Wu D., Yang S., Yao Q.A., Ye J., Yeh R.F., Zaveri J.S., Zhan M., Zhang G., Zhao Q., Zheng L., Zheng X.H., Zhong F.N., Zhong W., Zhou X., Zhu S., Zhu X., Smith H.O., Gibbs R.A., Myers E.W., Rubin G.M., Venter J.C. (2000). The genome sequence of *Drosophila melanogaster*. *Science* **287**: 2185-2195

Aizawa K., Akahira S., Akimura T., Arai A., Arakawa T., Carninci P., Hanagaki T., Hayatsu N., Hiraoka T., Hirozane T., Hodoyama Y., Imotani K., Ishii Y., Itoh M., Izawa M., Kawai J., Kojima Y., Konno H., Kusakabe M., Matsuyama T., Miyazaki A., Nakamura M., Nishi K., Nomura K., Numazaki R., Okazaki Y., Okido T., Owa C., Sakai C., Sakai K., Sasaki D., Sato K., Shibata K., Shibata Y., Shinagawa A., Shiraki T., Sogabe Y., Suzuki H., Tagawa A., Takahashi F., Tanaka T., Toya T., Watahiki A., Yamamura T., Yasunishi A., Yoshida K., Yoshiki A., Muramatsu M., Hayashizaki Y. (2000). RIKEN Mouse ESTs. *Unpublished*

Akao T., Kakehi Y., Itoh N., Ozdemir E., Shimizu T., Tachibana A., Sasaki M.S., Yoshida O. (1997). A high prevalence of functional inactivation by

methylation modification of p16INK4A/CDKN2/MTS1 gene in primary urothelial cancers. *Jpn J Cancer Res* **88**: 1078-1086

Akira S., Isshiki H., Sugita T., Tanabe O., Kinoshita S., Nishio Y., Nakajima T., Hirano T., Kishimoto T. (1990). A nuclear factor for IL-6 expression (NF-IL6) is a member of a C/EBP family. *Embo J* **9**: 1897-1906

Akiyama Y. (1998). TFSEARCH: Searching transcription factor binding sites v1.3 Parallel Application TRC Laboratory, Real World Computing Partnership, Japan. <http://www.cbrc.jp/research/db/TFSEARCH.html>

Alizadeh A.A., Ross D.T., Perou C.M., van de Rijn M. (2001). Towards a novel classification of human malignancies based on gene expression patterns. *J Pathol* **195**: 41-52

Altschul S.F., Gish W., Miller W., Myers E.W., Lipman D.J. (1990). Basic local alignment search tool. *J Mol Biol* **215**: 403-410 <http://www.ncbi.nlm.nih.gov/BLAST>

Ambrosio V., Borzacchiello G., Bruno F., Galati P., Roperto F. (2001). Uroplakin expression in the urothelial tumors of cows. *Vet Pathol* **38**: 657-660

Amiot M. (1990). Identification and analysis of cDNA clones encoding CD53. A pan-leukocyte antigen related to membrane transport proteins. *J Immunol* **145**: 4322-4325

Amonkar P., Murali G., Krishnamurthy S. (2001). Schistosoma induced squamous cell carcinoma of the bladder. *Indian J Pathol Microbiol* **44**: 363-364

Andrews P.W., Knowles B.B., Goodfellow P.N. (1981). A human cell-surface antigen defined by a monoclonal antibody and controlled by a gene on chromosome 12. *Somatic Cell Genet* **7**: 435-443

Angelisova P., Vlcek C., Stefanova I., Lipoldova M., Horejsi V. (1990). The human leucocyte surface antigen CD53 is a protein structurally similar to the CD37 and MRC OX-44 antigens. *Immunogenetics* **32**: 281-285

Bascom R.A., Connell G., Garcia-Heras J., Collins L., Ledbetter D., Molday R.S., Kalnins V., McInnes R.R. (1990). Molecular and ultrastructural characterization of the products of the human retinopathy candidate genes ROM1 and RDS. *Am. J. Hum. Genet.* **47 (suppl)**: A101

Bascom R.A., Manara S., Collins L., Molday R.S., Kalnins V.I., McInnes R.R. (1992). Cloning of the cDNA for a novel photoreceptor membrane protein (rom-1) identifies a disk rim protein family implicated in human retinopathies. *Neuron* **8**: 1171-1184

Bascom R.A., Schappert K., McInnes R.R. (1993). Cloning of the human and murine ROM1 genes: genomic organization and sequence conservation. *Hum Mol Genet* **2**: 385-391

- Baskin L.S., Hayward S.W., Sutherland R.A., DiSandro M.S., Thomson A.A. (1997). Cellular signaling in the bladder. *Front Biosci* **2**: d592-595
- Baskin L.S., Hayward S.W., Young P.F., Cunha G.R. (1996). Ontogeny of the rat bladder: smooth muscle and epithelial differentiation. *Acta Anat (Basel)* **155**: 163-171
- Baud E., Catilina P., Bignon Y.J. (1999). p16 involvement in primary bladder tumors: analysis of deletions and mutations. *Int J Oncol* **14**: 441-445
- Begy C., Bridges C.D. (1990). Nucleotide and predicted protein sequence of rat retinal degeneration slow (rds). *Nucleic Acids Res* **18**: 3058
- Bellacosa A., Lazo P.A., Bear S.E., Tsichlis P.N. (1991). The rat leukocyte antigen MRC OX-44 is a member of a new family of cell surface proteins which appear to be involved in growth regulation. *Mol Cell Biol* **11**: 2864-2872
- Bender C.M., Gonzalgo M.L., Gonzales F.A., Nguyen C.T., Robertson K.D., Jones P.A. (1999). Roles of cell division and gene transcription in the methylation of CpG islands. *Mol Cell Biol* **19**: 6690-6698
- Bender C.M., Zingg J.M., Jones P.A. (1998). DNA methylation as a target for drug design. *Pharm Res* **15**: 175-187
- Benedict W.F., Lerner S.P., Zhou J., Shen X., Tokunaga H., Czerniak B. (1999). Level of retinoblastoma protein expression correlates with p16 (MTS-1/INK4A/CDKN2) status in bladder cancer. *Oncogene* **18**: 1197-1203
- Benoit P., Gross M.S., Frachet P., Frezal J., Uzan G., Boucheix C., Nguyen V.C. (1991). Assignment of the human CD9 gene to chromosome 12 (region P13) by use of human specific DNA probes. *Hum Genet* **86**: 268-272
- Benson D.A., Karsch-Mizrachi I., Lipman D.J., Ostell J., Rapp B.A., Wheeler D.L. (2002). GenBank. *Nucleic Acids Res* **30**: 17-20 <http://www.ncbi.nlm.nih.gov>
- Berditchevski F. (2001). Complexes of tetraspanins with integrins: more than meets the eye. *J Cell Sci* **114**: 4143-4151
- Berditchevski F., Bazzoni G., Hemler M.E. (1995). Specific association of CD63 with the VLA-3 and VLA-6 integrins. *J Biol Chem* **270**: 17784-17790
- Berditchevski F., Zutter M.M., Hemler M.E. (1996). Characterization of novel complexes on the cell surface between integrins and proteins with 4 transmembrane domains (TM4 proteins). *Mol Biol Cell* **7**: 193-207
- Berggren P., Steineck G., Adolfsson J., Hansson J., Jansson O., Larsson P., Sandstedt B., Wijkstrom H., Hemminki K. (2001). p53 mutations in urinary bladder cancer. *Br J Cancer* **84**: 1505-1511
- Bernardin F., Yang Y., Cleaves R., Zahurak M., Cheng L., Civin C.I., Friedman A.D. (2002). TEL-AML1, expressed from t(12;21) in human acute lymphocytic leukemia, induces acute leukemia in mice. *Cancer Res* **62**: 3904-3908

- Bernardini S., Adessi G.L., Chezy E., Billerey C., Carbillot J.P., Bittard H. (2001). Influence of cigarette smoking on p53 gene mutations in bladder carcinomas. *Anticancer Res* **21**: 3001-3004
- Bestor T., Laudano A., Mattaliano R., Ingram V. (1988). Cloning and sequencing of a cDNA encoding DNA methyltransferase of mouse cells. The carboxyl-terminal domain of the mammalian enzymes is related to bacterial restriction methyltransferases. *J Mol Biol* **203**: 971-983
- Bianco T., Hussey D., Dobrovic A. (1999). Methylation-sensitive, single-strand conformation analysis (MS-SSCA): A rapid method to screen for and analyze methylation. *Hum Mutat* **14**: 289-293
- Bird A. (1992). The essentials of DNA methylation. *Cell* **70**: 5-8
- Bird A.P. (1986). CpG-rich islands and the function of DNA methylation. *Nature* **321**: 209-213
- Birling M.C., Tait S., Hardy R.J., Brophy P.J. (1999). A novel rat tetraspan protein in cells of the oligodendrocyte lineage. *J Neurochem* **73**: 2600-2608
- Boismenu R., Rhein M., Fischer W.H., Havran W.L. (1996). A role for CD81 in early T cell development. *Science* **271**: 198-200
- Bonaldo M.F., Lennon G., Soares M.B. (1996). Normalization and subtraction: two approaches to facilitate gene discovery. *Genome Res* **6**: 791-806
- Bonfils C., Beaulieu N., Chan E., Cotton-Montpetit J., MacLeod A.R. (2000). Characterization of the human DNA methyltransferase splice variant Dnmt1b. *J Biol Chem* **275**: 10754-10760
- Bornman D.M., Mathew S., Alsrude J., Herman J.G., Gabrielson E. (2001). Methylation of the E-cadherin gene in bladder neoplasia and in normal urothelial epithelium from elderly individuals. *Am J Pathol* **159**: 831-835
- Boucheix C., Rubinstein E. (2001). Tetraspanins. *Cell Mol Life Sci* **58**: 1189-1205
- Brakebusch C., Bouvard D., Stanchi F., Sakai T., Fassler R. (2002). Integrins in invasive growth. *J Clin Invest* **109**: 999-1006
- Brandau S., Bohle A. (2001). Bladder cancer. I. Molecular and genetic basis of carcinogenesis. *Eur Urol* **39**: 491-497
- Brooke G.P., Sopp P., Kwong L.S., Howard C.J. (1999). Molecular cloning of cattle CD63 and evidence for high level expression on subpopulations of dendritic cells. *Immunogenetics* **49**: 812-814
- Bucher P. (1990). Weight matrix descriptions of four eukaryotic RNA polymerase II promoter elements derived from 502 unrelated promoter sequences. *J Mol Biol* **212**: 563-578

- Burchert A., Notter M., Dietrich Menssen H., Schwartz S., Knauf W., Neubauer A., Thiel E. (1999). CD82 (KAI1), a member of the tetraspan family, is expressed on early haemopoietic progenitor cells and up-regulated in distinct human leukaemias. *Br J Haematol* **107**: 494-504
- Burkitt G. (1996). Wheater's Basic Histopathology. Churchill Livingstone, Inc., New York, NY, USA.
- Cairns P., Polascik T.J., Eby Y., Tokino K., Califano J., Merlo A., Mao L., Herath J., Jenkins R., Westra W., Rutter J.L., Buckler A., Gavielson E., Tockman M., Cho K.R., Hedrick L., Bova G.S., Issacs W., Schwab D., Sidransky D. (1995). Frequency of homozygous deletion at p16/CDKN2 in primary human tumours. *Nat Genet* **11**: 210-212
- Castellvi-Bel S., Mila M. (2001). Genes responsible for nonspecific mental retardation. *Mol Genet Metab* **72**: 104-108
- Chang A., Hammond T.G., Sun T.T., Zeidel M.L. (1994). Permeability properties of the mammalian bladder apical membrane. *Am J Physiol* **267**: C1483-1492
- Charache S., Dover G., Smith K., Talbot C.C., Jr., Moyer M., Boyer S. (1983). Treatment of sickle cell anemia with 5-azacytidine results in increased fetal hemoglobin production and is associated with nonrandom hypomethylation of DNA around the gamma-delta-beta-globin gene complex. *Proc Natl Acad Sci U S A* **80**: 4842-4846
- Chlapowski F.J., Bonneville M.A., Staehelin L.A. (1972). Lumenal plasma membrane of the urinary bladder. II. Isolation and structure of membrane components. *J Cell Biol* **53**: 92-104
- Cho S.K., Kwon D.N., Chu H.J., Kim J.H. (1999). Cloning of rabbit uroplakin 1b cDNA.
- Chretien S., Dubart A., Beaupain D., Raich N., Grandchamp B., Rosa J., Goossens M., Romeo P.H. (1988). Alternative transcription and splicing of the human porphobilinogen deaminase gene result either in tissue-specific or in housekeeping expression. *Proc Natl Acad Sci U S A* **85**: 6-10
- Claas C., Seiter S., Claas A., Savelyeva L., Schwab M., Zoller M. (1998). Association between the rat homologue of CO-029, a metastasis-associated tetraspanin molecule and consumption coagulopathy. *J Cell Biol* **141**: 267-280
- Clark D.A., Coker R. (1998). Transforming growth factor-beta (TGF-beta). *Int J Biochem Cell Biol* **30**: 293-298
- Clark S.J., Frommer M. (1997). Bisulphite genomic sequencing of methylated cytosines. In "Laboratory methods for the detection of mutations and polymorphisms in DNA" (Taylor G.R., Ed.), pp. 151-162, CRC Press, Inc., Boca Raton, FL, USA.

- Clark S.J., Harrison J., Molloy P.L. (1997). Sp1 binding is inhibited by (m)Cp(m)CpG methylation. *Gene* **195**: 67-71
- Clifton S., Johnson S.L., Blumberg B., Song J., Hillier L., Pape D., Martin J., Wylie T., Underwood K., Theising B., Bowers Y., Person B., Gibbons M., Harvey N., Ritter E., Jackson Y., McCann R., Waterston R., Wilson R. (1999). WashU Xenopus EST project, 1999. *GenBank direct submission*
- Consortium (1998). Genome sequence of the nematode *C. elegans*: a platform for investigating biology. The *C. elegans* Sequencing Consortium. *Science* **282**: 2012-2018
- Courseaux A., Grosgeorge J., Gaudray P., Pannett A.A.J., Forbes S.A., Williamson C., Bassett D., Thakker R.V., Teh B.T., Farnebo F., Shepherd J., Skogseid B., Larsson C., Giraud S., Zhang C.X., Salandre J., Calender A. (1996). Definition of the Minimal MEN1 Candidate Area Based on a 5-Mb Integrated Map of Proximal 11q13. *Genomics* **37**: 345-353
- Covey J.M., Zaharko D.S. (1984). Effects of dose and duration of exposure on 5-aza-2'-deoxycytidine cytotoxicity for L1210 leukemia in vitro. *Cancer Treat Rep* **68**: 1475-1481
- Crotta S., Still A., Wack A., D'Andrea A., Nuti S., D'Oro U., Mosca M., Filliponi F., Brunetto R.M., Bonino F., Abrignani S., Valiante N.M. (2002). Inhibition of natural killer cells through engagement of CD81 by the major hepatitis C virus envelope protein. *J Exp Med* **195**: 35-41
- Dahia P.L., Aguiar R.C., Honegger J., Fahlbush R., Jordan S., Lowe D.G., Lu X., Clayton R.N., Besser G.M., Grossman A.B. (1998). Mutation and expression analysis of the p27/kip1 gene in corticotrophin-secreting tumours. *Oncogene* **16**: 69-76
- Dalbagni G., Presti J., Reuter V., Fair W.R., Cordon-Cardo C. (1993a). Genetic alterations in bladder cancer. *Lancet* **342**: 469-471
- Dalbagni G., Presti J.C., Jr., Reuter V.E., Zhang Z.F., Sarkis A.S., Fair W.R., Cordon-Cardo C. (1993b). Molecular genetic alterations of chromosome 17 and p53 nuclear overexpression in human bladder cancer. *Diagn Mol Pathol* **2**: 4-13
- Davern K.M., Wright M.D., Herrmann V.R., Mitchell G.F. (1991). Further characterisation of the *Schistosoma japonicum* protein Sj23, a target antigen of an immunodiagnostic monoclonal antibody. *Mol Biochem Parasitol* **48**: 67-75
- DeRisi J., Penland L., Brown P.O., Bittner M.L., Meltzer P.S., Ray M., Chen Y., Su Y.A., Trent J.M. (1996). Use of a cDNA microarray to analyse gene expression patterns in human cancer. *Nat Genet* **14**: 457-460
- Domanico S.Z., Pelletier A.J., Havran W.L., Quaranta V. (1997). Integrin alpha 6A beta 1 induces CD81-dependent cell motility without engaging the extracellular matrix migration substrate. *Mol Biol Cell* **8**: 2253-2265

- Dong J.T., Lamb P.W., Rinker-Schaeffer C.W., Vukanovic J., Ichikawa T., Isaacs J.T., Barrett J.C. (1995). KAI1, a metastasis suppressor gene for prostate cancer on human chromosome 11p11.2. *Science* **268**: 884-886
- Douglas G.C., Hossler F.E. (1995). Vascular anatomy of the rabbit ureter. *Anat Rec* **242**: 47-56
- Dracopoli N.C., Haines J.L., Korf B.R., Moir D.T., Morton C.C., Seidman C.E., Seidman J.G., Smith D.R. (1994). Current protocols in human genetics. John Wiley & Sons Inc., New York, NY, USA.
- Dynan W.S., Tjian R. (1983). The promoter-specific transcription factor Sp1 binds to upstream sequences in the SV40 early promoter. *Cell* **35**: 79-87
- Eichelberg D., Wessing A. (1971). Electron microscopic studies of the renal tubules (Malpighian tubules) in *Drosophila melanogaster*. II. Transcellular membrane-bound transport. *Z Zellforsch Mikrosk Anat* **121**: 127-152
- Elkahloun A.G., Krizman D.B., Wang Z., Hofmann T.A., Roe B., Meltzer P.S. (1997). Transcript mapping in a 46-kb sequenced region at the core of 12q13.3 amplification in human cancers. *Genomics* **42**: 295-301
- Emi N., Kitaori K., Seto M., Ueda R., Saito H., Takahashi T. (1993). Isolation of a novel cDNA clone showing marked similarity to ME491/CD63 superfamily. *Immunogenetics* **37**: 193-198
- Engels B. (1992). Amplify v1.2 University of Wisconsin, Madison, WI, USA.
- Esrig D., Elmajian D., Groshen S., Freeman J.A., Stein J.P., Chen S.C., Nichols P.W., Skinner D.G., Jones P.A., Cote R.J. (1994). Accumulation of nuclear p53 and tumor progression in bladder cancer. *N Engl J Med* **331**: 1259-1264
- Falls J.G., Pulford D.J., Wylie A.A., Jirtle R.L. (1999). Genomic imprinting: implications for human disease. *Am J Pathol* **154**: 635-647
- Fan J., Brindley P.J. (1998). Characterization of cDNAs encoding a new family of tetraspanins from schistosomes--the Sj25 family. *Gene* **219**: 1-8
http://www.elsevier.com:80/cgi-bin/cas/tree/store/gene/cas_sub/browse/browse.cgi?year=1998&volume=219&issue=1-2&aid=11520
- Fan J., Hooker C.W., McManus D.P., Brindley P.J. (1997). A new member of the transmembrane 4 superfamily (TM4SF) of proteins from schistosomes, expressed by larval and adult *Schistosoma japonicum*. *Biochim Biophys Acta* **1329**: 18-25
- Felsenstein J. (1993a). CONSENSE: Consensus tree program v3.5c University of Washington, Seattle, WA, USA. <http://evolution.genetics.washington.edu/phylip.html>
- Felsenstein J. (1993b). DNAPARS: DNA parsimony program v3.5c University of Washington, Seattle, WA, USA. <http://evolution.genetics.washington.edu/phylip.html>

Felsenstein J. (1993c). SEQBOOT: Bootstrap, jackknife, or permutation resampling of molecular sequence, restriction site, gene frequency or character data v3.5c University of Washington, Seattle, WA, USA. <http://evolution.genetics.washington.edu/phylip.html>

Ferguson A.T., Vertino P.M., Spitzner J.R., Baylin S.B., Muller M.T., Davidson N.E. (1997). Role of estrogen receptor gene demethylation and DNA methyltransferase.DNA adduct formation in 5-aza-2'deoxyctydine-induced cytotoxicity in human breast cancer cells. *J Biol Chem* **272**: 32260-32266

Finch J.L. (1998). Cloning and characterisation of the human uroplakin 1B gene. Department of Surgery, The University of Adelaide, Adelaide, Australia.

Finch J.L., Miller J., Aspinall J.O., Cowled P.A. (1999). Cloning of the human uroplakin 1B cDNA and analysis of its expression in urothelial-tumor cell lines and bladder-carcinoma tissue. *Int J Cancer* **80**: 533-538

Finch J.L., Webb G.C., Evdokiou A., Cowled P.A. (1997). Chromosomal localization of the human urothelial "tetraspan" gene, UPK1B, to 3q13.3-q21 and detection of a TaqI polymorphism. *Genomics* **40**: 501-503

Fitter S., Seldin M.F., Ashman L.K. (1998). Characterisation of the mouse homologue of CD151 (PETA-3/SFA-1); genomic structure, chromosomal localisation and identification of 2 novel splice forms. *Biochim Biophys Acta* **1398**: 75-85 http://www.elsevier.com:80/cgi-bin/cas/tree/store/bbaexp/cas_sub/browse/browse.cgi?year=1998&volume=1398&issue=1&aid=93102

Fitter S., Tetaz T.J., Berndt M.C., Ashman L.K. (1995). Molecular cloning of cDNA encoding a novel platelet-endothelial cell tetra-span antigen, PETA-3. *Blood* **86**: 1348-1355

Fowler J.E., Jr., Mariano M., Lau J.L. (1987). Interaction of urinary Tamm-Horsfall protein with transitional cells and transitional epithelium. *J Urol* **138**: 446-448

Friedrich M.G., Blind C., Milde-Langosch K., Erbersdobler A., Conrad S., Loning T., Hammerer P., Huland H. (2001). Frequent p16/MTS1 inactivation in early stages of urothelial carcinoma of the bladder is not associated with tumor recurrence. *Eur Urol* **40**: 518-524

Frommer M., McDonald L.E., Millar D.S., Collis C.M., Watt F., Grigg G.W., Molloy P.L., Paul C.L. (1992). A genomic sequencing protocol that yields a positive display of 5-methylcytosine residues in individual DNA strands. *Proc Natl Acad Sci U S A* **89**: 1827-1831

Fujimoto K., Yamada Y., Okajima E., Kakizoe T., Sasaki H., Sugimura T., Terada M. (1992). Frequent association of p53 gene mutation in invasive bladder cancer. *Cancer Res* **52**: 1393-1398

Gabriel J.M., Higgins M.J., Gebuhr T.C., Shows T.B., Saitoh S., Nicholls R.D. (1998). A model system to study genomic imprinting of human genes. *Proc Natl Acad Sci U S A* **95**: 14857-14862

- Geisert E.E., Jr., Murphy T.P., Irwin M.H., Larjava H. (1991). A novel cell adhesion molecule, G-CAM, found on cultured rat glia. *Neurosci Lett* **133**: 262-266
- Geisert E.E., Jr., Yang L., Irwin M.H. (1996). Astrocyte growth, reactivity, and the target of the antiproliferative antibody, TAPA. *J Neurosci* **16**: 5478-5487
- Gelfand M., Weinberg R.W., Castle W.M. (1967). Relation between carcinoma of the bladder and infestation with Schistosoma haematobium. *Lancet* **1**: 1249-1251
- Gil M.L., Vita N., Lebel-Binay S., Miloux B., Chalon P., Kaghad M., Marchiol-Fournigault C., Conjeaud H., Caput D., Ferrara P. (1992). A member of the tetraspan transmembrane protein superfamily is recognized by a monoclonal antibody raised against an HLA class I-deficient, lymphokine-activated killer-susceptible, B lymphocyte line. Cloning and preliminary functional studies. *J Immunol* **148**: 2826-2833
- Glazer R.I., Knodel M.C. (1984). 1-beta-D-arabinosyl-5-azacytosine. Cytocidal activity and effects on the synthesis and methylation of DNA in human colon carcinoma cells. *Mol Pharmacol* **26**: 381-387
- Gonzalez-Zulueta M., Bender C.M., Yang A.S., Nguyen T., Beart R.W., Van Tornout J.M., Jones P.A. (1995). Methylation of the 5' CpG island of the p16/CDKN2 tumor suppressor gene in normal and transformed human tissues correlates with gene silencing. *Cancer Res* **55**: 4531-4535
- Gonzalgo M.L., Jones P.A. (1997). Rapid quantitation of methylation differences at specific sites using methylation-sensitive single nucleotide primer extension (Ms-SNuPE). *Nucleic Acids Res* **25**: 2529-2531
- Gorin M.B., Snyder S., To A., Narfstrom K., Curtis R. (1993). The cat RDS transcript: candidate gene analysis and phylogenetic sequence analysis. *Mamm Genome* **4**: 544-548
- Greger V., Passarge E., Hopping W., Messmer E., Horsthemke B. (1989). Epigenetic changes may contribute to the formation and spontaneous regression of retinoblastoma. *Hum Genet* **83**: 155-158
- Griffiths M.M., Encinas J.A., Remmers E.F., Kuchroo V.K., Wilder R.L. (1999). Mapping autoimmunity genes. *Curr Opin Immunol* **11**: 689-700
- Gulko P.S., Kawahito Y., Remmers E.F., Reese V.R., Wang J., Dracheva S.V., Ge L., Longman R.E., Shepard J.S., Cannon G.W., Sawitzke A.D., Wilder R.L., Griffiths M.M. (1998). Identification of a new non-major histocompatibility complex genetic locus on chromosome 2 that controls disease severity in collagen-induced arthritis in rats. *Arthritis Rheum* **41**: 2122-2131
- Guo X.Z., Friess H., Di Mola F.F., Heinicke J.M., Abou-Shady M., Gruber H.U., Baer H.U., Zimmermann A., Korc M., Buchler M.W. (1998). KAI1, a new metastasis suppressor gene, is reduced in metastatic hepatocellular carcinoma. *Hepatology* **28**: 1481-1488

- Guo X.Z., Friess H., Shao X.D., Liu M.P., Xia Y.T., Xu J.H., Buchler M.W. (2000). KAI1 gene is differently expressed in papillary and pancreatic cancer: influence on metastasis. *World J Gastroenterol* **6**: 866-871
- Hall T.A. (2001). BioEdit v5.0.9 North Carolina State University, Raleigh, NC, USA.
- Harano H., Wang C., Gao J., Uchida T. (1999). p53 tumor suppressor gene mutation and prognosis in 105 cases of bladder cancer--the relationship between mutation of the p53 gene with clinicopathological features and smoking. *Nippon Hinyokika Gakkai Zasshi* **90**: 487-495
- Hartge P., Harvey E.B., Linehan W.M., Silverman D.T., Sullivan J.W., Hoover R.N., Fraumeni J.F., Jr. (1990). Unexplained excess risk of bladder cancer in men. *J Natl Cancer Inst* **82**: 1636-1640
- Hartmann A., Moser K., Kriegmair M., Hofstetter A., Hofstaedter F., Knuechel R. (1999). Frequent genetic alterations in simple urothelial hyperplasias of the bladder in patients with papillary urothelial carcinoma. *Am J Pathol* **154**: 721-727
- Hartmann A., Schlake G., Zaak D., Hungerhuber E., Hofstetter A., Hofstaedter F., Knuechel R. (2002). Occurrence of chromosome 9 and p53 alterations in multifocal dysplasia and carcinoma in situ of human urinary bladder. *Cancer Res* **62**: 809-818
- Heby O. (1995). DNA methylation and polyamines in embryonic development and cancer. *Int J Dev Biol* **39**: 737-757
- Heinemeyer T., Chen X., Karas H., Kel A.E., Kel O.V., Liebich I., Meinhardt T., Reuter I., Schacherer F., Wingender E. (1999). Expanding the TRANSFAC database towards an expert system of regulatory molecular mechanisms. *Nucleic Acids Res* **27**: 318-322
- Heinemeyer T., Wingender E., Reuter I., Hermjakob H., Kel A.E., Kel O.V., Ignatieva E.V., Ananko E.A., Podkolodnaya O.A., Kolpakov F.A., Podkolodny N.L., Kolchanov N.A. (1998). Databases on transcriptional regulation: TRANSFAC, TRRD and COMPEL. *Nucleic Acids Res* **26**: 362-367
- Hemler M.E. (2001). Specific tetraspanin functions. *J Cell Biol* **155**: 1103-1107
- Hemler M.E., Mannion B.A., Berditchevski F. (1996). Association of TM4SF proteins with integrins: relevance to cancer. *Biochim Biophys Acta* **1287**: 67-71
- Herman J.G., Baylin S.B. (1994). Methylation Specific PCR. In "Current protocols in human genetics" (Dracopoli N.C., Haines J.L., Korf B.R., Moir D.T., Morton C.C., Seidman C.E., Seidman J.G., Smith D.R., Eds.), John Wiley & Sons Inc., New York, NY, USA.
- Herman J.G., Graff J.R., Myohanen S., Nelkin B.D., Baylin S.B. (1996). Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. *Proc Natl Acad Sci U S A* **93**: 9821-9826

- Herman J.G., Merlo A., Mao L., Lapidus R.G., Issa J.P., Davidson N.E., Sidransky D., Baylin S.B. (1995). Inactivation of the CDKN2/p16/MTS1 gene is frequently associated with aberrant DNA methylation in all common human cancers. *Cancer Res* **55**: 4525-4530
- Hicks R.M. (1965). The fine structure of the transitional epithelium of rat ureter. *J Cell Biol* **26**: 25-48
- Hicks R.M., Ketterer B. (1969). Hexagonal lattice of subunits in the thick luminal membrane of the rat urinary bladder. *Nature* **224**: 1304-1305
- Higashiyama M., Doi O., Kodama K., Yokouchi H., Adachi M., Huang C.L., Taki T., Kasugai T., Ishiguro S., Nakamori S., Miyake M. (1997). Immunohistochemically detected expression of motility-related protein-1 (MRP-1/CD9) in lung adenocarcinoma and its relation to prognosis. *Int J Cancer* **74**: 205-211
- Higashiyama M., Kodama K., Yokouchi H., Takami K., Adachi M., Taki T., Ishiguro S., Nakamori S., Yoshie O., Miyake M. (1998). KAI1/CD82 expression in nonsmall cell lung carcinoma is a novel, favorable prognostic factor: an immunohistochemical analysis. *Cancer* **83**: 466-474
- Ho A., Dowdy S.F. (2002). Regulation of G(1) cell-cycle progression by oncogenes and tumor suppressor genes. *Curr Opin Genet Dev* **12**: 47-52
- Hogan T.F., Lamm D.L., Riggs D.R., Ducatman A.M. (2001). Smoking and bladder cancer. *W V Med J* **97**: 29-31
- Hoodless P.A., Wrana J.L. (1998). Mechanism and function of signaling by the TGF beta superfamily. *Curr Top Microbiol Immunol* **228**: 235-272
- Horejsi V., Vlcek C. (1991). Novel structurally distinct family of leucocyte surface glycoproteins including CD9, CD37, CD53 and CD63. *FEBS Lett* **288**: 1-4
- Horvath G., Serru V., Clay D., Billard M., Boucheix C., Rubinstein E. (1998). CD19 is linked to the integrin-associated tetraspans CD9, CD81, and CD82. *J Biol Chem* **273**: 30537-30543 <http://www.jbc.org/cgi/content/full/273/46/30537>
- Hosokawa Y., Ueyama E., Morikawa Y., Maeda Y., Seto M., Senba E. (1999). Molecular cloning of a cDNA encoding mouse A15, a member of the transmembrane 4 superfamily, and its preferential expression in brain neurons. *Neurosci Res* **35**: 281-290
- Hotta H., Hara I., Miyamoto H., Homma M. (1991). Overexpression of the human melanoma-associated antigen ME491 partially suppresses in vivo malignant phenotypes of H-ras-transformed NIH3T3 cells in athymic nude mice. *Melanoma Res* **1**: 125-132
- Hotta H., Miyamoto H., Hara I., Takahashi N., Homma M. (1992). Genomic structure of the ME491/CD63 antigen gene and functional analysis of the 5'-flanking regulatory sequences. *Biochem Biophys Res Commun* **185**: 436-442

- Hotta H., Ross A.H., Huebner K., Isobe M., Wendeborn S., Chao M.V., Ricciardi R.P., Tsujimoto Y., Croce C.M., Koprowski H. (1988). Molecular cloning and characterization of an antigen associated with early stages of melanoma tumor progression. *Cancer Res* **48**: 2955-2962
- Hotta H., Takahashi N., Homma M. (1989). Transcriptional enhancement of the human gene encoding for a melanoma-associated antigen (ME491) in association with malignant transformation. *Jpn J Cancer Res* **80**: 1186-1191
- Hu R.J., Lee M.P., Connors T.D., Johnson L.A., Burn T.C., Su K., Landes G.M., Feinberg A.P. (1997). A 2.5-Mb transcript map of a tumor-suppressing subchromosomal transferable fragment from 11p15.5, and isolation and sequence analysis of three novel genes. *Genomics* **46**: 9-17
- Huang C.I., Kohno N., Ogawa E., Adachi M., Taki T., Miyake M. (1998). Correlation of reduction in MRP-1/CD9 and KAI1/CD82 expression with recurrences in breast cancer patients. *Am J Pathol* **153**: 973-983
- Ichikawa T., Ichikawa Y., Dong J., Hawkins A.L., Griffin C.A., Isaacs W.B., Oshimura M., Barrett J.C., Isaacs J.T. (1992). Localization of metastasis suppressor gene(s) for prostatic cancer to the short arm of human chromosome 11. *Cancer Res* **52**: 3486-3490
- Ikeyama S., Koyama M., Yamaoko M., Sasada R., Miyake M. (1993). Suppression of cell motility and metastasis by transfection with human motility-related protein (MRP-1/CD9) DNA. *J Exp Med* **177**: 1231-1237
- Imai T., Fukudome K., Takagi S., Nagira M., Furuse M., Fukuhara N., Nishimura M., Hinuma Y., Yoshie O. (1992). C33 antigen recognized by monoclonal antibodies inhibitory to human T cell leukemia virus type 1-induced syncytium formation is a member of a new family of transmembrane proteins including CD9, CD37, CD53, and CD63. *J Immunol* **149**: 2879-2886
- Inal J., Bickle Q. (1995). Sequence and immunogenicity of the 23-kDa transmembrane antigen of Schistosoma haematobium. *Mol Biochem Parasitol* **74**: 217-221
- Irwin M.H., Geisert E.E., Jr. (1993). The upregulation of a glial cell surface antigen at the astrocytic scar in the rat. *Neurosci Lett* **154**: 57-60
- Ishikawa J., Xu H.J., Hu S.X., Yandell D.W., Maeda S., Kamidono S., Benedict W.F., Takahashi R. (1991). Inactivation of the retinoblastoma gene in human bladder and renal cell carcinomas. *Cancer Res* **51**: 5736-5743
- Jackson P., Grimm M.O., Kingsley E.A., Brosius U., Antalis T., Yardley G., Russell P.J. (2002). Relationship between expression of KAI1 metastasis suppressor gene, mRNA levels and p53 in human bladder and prostate cancer cell lines. *Urol Oncol* **7**: 99-104
- Jackson P., Kingsley E.A., Russell P.J. (2000a). Inverse correlation between KAI1 mRNA levels and invasive behaviour in bladder cancer cell lines. *Cancer Lett* **156**: 9-17

- Jackson P., Millar D., Kingsley E., Yardley G., Ow K., Clark S., Russell P.J. (2000b). Methylation of a CpG island within the promoter region of the KAI1 metastasis suppressor gene is not responsible for down-regulation of KAI1 expression in invasive cancers or cancer cell lines. *Cancer Lett* **157**: 169-176
- Jackson P., Puisieux A. (2000). Is the KAI1 metastasis suppressor gene a cellular target of p53? A review of current evidence. *Biochem Biophys Res Commun* **278**: 499-502
- Jankowski S.A., De Jong P., Meltzer P.S. (1995). Genomic structure of SAS, a member of the transmembrane 4 superfamily amplified in human sarcomas. *Genomics* **25**: 501-506
- Jones P.A. (1985a). Altering gene expression with 5-azacytidine. *Cell* **40**: 485-486
- Jones P.A. (1985b). Effects of 5-azacytidine and its 2'-deoxyderivative on cell differentiation and DNA methylation. *Pharmacol Ther* **28**: 17-27
- Jones P.A., Taylor S.M. (1980). Cellular differentiation, cytidine analogs and DNA methylation. *Cell* **20**: 85-93
- Jones P.A., Taylor S.M., Wilson V.L. (1983). Inhibition of DNA methylation by 5-azacytidine. *Recent Results Cancer Res* **84**: 202-211
- Jüttermann R., Li E., Jaenisch R. (1994). Toxicity of 5-aza-2'-deoxycytidine to mammalian cells is mediated primarily by covalent trapping of DNA methyltransferase rather than DNA demethylation. *Proc Natl Acad Sci U S A* **91**: 11797-11801
- Kageyama S., Yoshiki T., Isono T., Tanaka T., Kim C.J., Yuasa T., Okada Y. (2002). High expression of human uroplakin ia in urinary bladder transitional cell carcinoma. *Jpn J Cancer Res* **93**: 523-531
- Kaghad M., Bonnet H., Yang A., Creancier L., Biscan J.C., Valent A., Minty A., Chalon P., Lelias J.M., Dumont X., Ferrara P., McKeon F., Caput D. (1997). Monoallelically expressed gene related to p53 at 1p36, a region frequently deleted in neuroblastoma and other human cancers. *Cell* **90**: 809-819
- Kaji K., Oda S., Shikano T., Ohnuki T., Uematsu Y., Sakagami J., Tada N., Miyazaki S., Kudo A. (2000). The gamete fusion process is defective in eggs of Cd9-deficient mice. *Nat Genet* **24**: 279-282
- Kallin B., de Martin R., Etzold T., Sorrentino V., Philipson L. (1991). Cloning of a growth arrest-specific and transforming growth factor beta-regulated gene, TI 1, from an epithelial cell line. *Mol Cell Biol* **11**: 5338-5345
- Kamb A., Gruis N.A., Weaver-Feldhaus J., Liu Q., Harshman K., Tavtigian S.V., Stockert E., Day R.S., 3rd, Johnson B.E., Skolnick M.H. (1994). A cell cycle regulator potentially involved in genesis of many tumor types. *Science* **264**: 436-440

- Kanerva J., Niini T., Vettentranta K., Riikonen P., Makipernaa A., Karhu R., Knuutila S., Saarinen-Pihkala U.M. (2001). Loss at 12p detected by comparative genomic hybridization (CGH): association with TEL-AML1 fusion and favorable prognostic features in childhood acute lymphoblastic leukemia (ALL). A multi-institutional study. *Med Pediatr Oncol* **37**: 419-425
- Kaprielian Z., Cho K.O., Hadjigyrou M., Patterson P.H. (1995). CD9, a major platelet cell surface glycoprotein, is a ROCA antigen and is expressed in the nervous system. *J Neurosci* **15**: 562-573
- Kardong K.V. (1995). Vertebrates: comparative anatomy, function, evolution. Wm. C. Brown Publishers, Dubuque, IA, USA.
- Karpf A.R., Jones D.A. (2002). Reactivating the expression of methylation silenced genes in human cancer. *Oncogene* **21**: 5496-5503
- Kaufmann O., Volmerig J., Dietel M. (2000). Uroplakin III is a highly specific and moderately sensitive immunohistochemical marker for primary and metastatic urothelial carcinomas. *Am J Clin Pathol* **113**: 683-687
- Kedzierski W., Moghrabi W.N., Allen A.C., Jablonski-Stiemke M.M., Azarian S.M., Bok D., Travis G.H. (1996). Three homologs of rds/peripherin in Xenopus laevis photoreceptors that exhibit covalent and non-covalent interactions. *J Cell Sci* **109**: 2551-2560
- Kedzierski W., Weng J., Travis G.H. (1999). Analysis of the rds/peripherin.rom1 complex in transgenic photoreceptors that express a chimeric protein. *J Biol Chem* **274**: 29181-29187
- Kerr L.D., Miller D.B., Matrisian L.M. (1990). TGF-beta 1 inhibition of transin/stromelysin gene expression is mediated through a Fos binding sequence. *Cell* **61**: 267-278
- Knudson A.G., Jr., Strong L.C., Anderson D.E. (1973). Heredity and cancer in man. *Prog Med Genet* **9**: 113-158
- Kobayashi T., Mekada E. (1999). *GenBank direct submission*
- Kolch W., Kotwaliwale A., Vass K., Jonosch P. (2002). The role of Raf kinases in malignant transformation. *Exp. Rev. Mol. Med.* **25 April** <http://www.expertreviews.org/02004386h.htm>
- Konno H., Aizawa K., Akahira S., Akiyama J., Arakawa T., Carninci P., Endo T., Fukuda S., Fukunishi Y., Hara A., Hayatsu N., Hirozane T., Hori F., Ishii Y., Ishikawa J., Ishikawa T., Itoh M., Izawa M., Kadota K., Kagawa I., Kai C., Kawai J., Kikuchi N., Kiyosawa H., Kojima Y., Kondo S., Koya S., Kurihara C., Kusakabe M., Matsuyama T., Miki R., Mizuno Y., Nakamura M., Oda H., Okazaki Y., Ono T., Owa C., Saito H., Sakai C., Sato K., Shibata K., Shibata Y., Shigemoto Y., Shinagawa A., Shiraki T., Sogabe Y., Sugahara Y., Suzuki H., Suzuki H., Tagawa A., Takahashi F., Tominaga N., Toya T., Tsunoda Y., Watahiki A., Watanabe S., Yamamura T., Yamanaka I., Yano R., Yasunishi A.,

- Yokota T., Yoshida K., Yoshiki A., Yoshino M., Muramatsu M., Hayashizaki Y. (2000). RIKEN Mouse ESTs. *Unpublished*
- Kopczynski C.C., Davis G.W., Goodman C.S. (1996). A neural tetraspanin, encoded by late bloomer, that facilitates synapse formation. *Science* **271**: 1867-1870
- Kortschak R.D. (1998). AddGaps The University of Adelaide, Adelaide.
- Koyama Y., Suzuki M., Yoshida T. (1998). CD63, a member of tetraspan transmembrane protein family, induces cellular spreading by reaction with monoclonal antibody on substrata. *Biochem Biophys Res Commun* **246**: 841-846
- Koyama Y., Yamanoha B., Yoshida T. (1990). A novel monoclonal antibody induces the differentiation of monocyte leukemic cells. *Biochem Biophys Res Commun* **168**: 898-904
- Kwon D.N., Chu H.J., Cho S.K., Kim J.H. (1999). Cloning of pig uroplakin Ia cDNA. *GenBank direct submission*
- Larget B., Simon D. (1999). Markov chain Monte Carlo algorithms for the Bayesian analysis of phylogenetic trees. *Molecular Biology and Evolution* **16**: 750-759
- Le Naour F., Prenant M., Francastel C., Rubinstein E., Uzan G., Boucheix C. (1996). Transcriptional regulation of the human CD9 gene: characterization of the 5'-flanking region. *Oncogene* **13**: 481-486
- Le Naour F., Rubinstein E., Jasmin C., Prenant M., Boucheix C. (2000). Severely reduced female fertility in CD9-deficient mice. *Science* **287**: 319-321
- Lebel-Binay S., Lagaudriere C., Fradelizi D., Conjeaud H. (1995). CD82, member of the tetra-span-transmembrane protein family, is a costimulatory protein for T cell activation. *J Immunol* **155**: 101-110
- Lee K.W., Shalaby K.A., Medhat A.M., Shi H., Yang Q., Karim A.M., LoVerde P.T. (1995). Schistosoma mansoni: characterization of the gene encoding Sm23, an integral membrane protein. *Exp Parasitol* **80**: 155-158
- Levy S., Kuo C.C. (1998). *GenBank direct submission*
- Lewis S.A. (2000). Everything you wanted to know about the bladder epithelium but were afraid to ask. *Am J Physiol Renal Physiol* **278**: F867-874
- Li E., Beard C., Jaenisch R. (1993a). Role for DNA methylation in genomic imprinting. *Nature* **366**: 362-365
- Li E., Bestor T.H., Jaenisch R. (1992). Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell* **69**: 915-926
- Li S.H., McInnis M.G., Margolis R.L., Antonarakis S.E., Ross C.A. (1993b). Novel triplet repeat containing genes in human brain: cloning, expression, and length polymorphisms. *Genomics* **16**: 572-579

- Li S.M., Zhang Z.T., Chan S., McLenan O., Dixon C., Taneja S., Lepor H., Sun T.T., Wu X.R. (1999). Detection of circulating uroplakin-positive cells in patients with transitional cell carcinoma of the bladder. *J Urol* **162**: 931-935
- Liang G., Gonzales F.A., Jones P.A., Orntoft T.F., Thykjaer T. (2002). Analysis of gene induction in human fibroblasts and bladder cancer cells exposed to the methylation inhibitor 5-aza-2'-deoxycytidine. *Cancer Res* **62**: 961-966
- Lin J.H., Zhao H., Sun T.T. (1995). A tissue-specific promoter that can drive a foreign gene to express in the suprabasal urothelial cells of transgenic mice. *Proc Natl Acad Sci U S A* **92**: 679-683
- Lobban E.D., Smith B.A., Hall G.D., Harnden P., Roberts P., Selby P.J., Trejdosiewicz L.K., Southgate J. (1998). Uroplakin gene expression by normal and neoplastic human urothelium. *Am J Pathol* **153**: 1957-1967
- Logothetis C., Swanson D., Amato R., Banks M., Finn L., Ayala A., Ro J., Babaian R., Dinney C., Ellerhorst J., Hall C., von Eschenbach A. (1996). Optimal delivery of perioperative chemotherapy: preliminary results of a randomized, prospective, comparative trial of preoperative and postoperative chemotherapy for invasive bladder carcinoma. *J Urol* **155**: 1241-1245
- Lombardi D.P., Geraerts J., Foley J.F., Chiao C., Lamb P.W., Barrett J.C. (1999). Loss of KAI1 expression in the progression of colorectal cancer. *Cancer Res* **59**: 5724-5731
- Lopez R., Lloyd A. (1994). ClustalW WWW Service at the European Bioinformatics Institute The ClustalWWW server at the EBIembnet.news volume 4.2 1997
http://www2.ebi.ac.uk/embnet.news/vol4_3/clustalw1.html.
<http://www.ebi.ac.uk/clustalw>
- Lu J.J., Kakehi Y., Takahashi T., Wu X.X., Yuasa T., Yoshiki T., Okada Y., Terachi T., Ogawa O. (2000). Detection of circulating cancer cells by reverse transcription-polymerase chain reaction for uroplakin II in peripheral blood of patients with urothelial cancer. *Clin Cancer Res* **6**: 3166-3171
- Ma J., Norton J.C., Allen A.C., Burns J.B., Hasel K.W., Burns J.L., Sutcliffe J.G., Travis G.H. (1995). Retinal degeneration slow (rds) in mouse results from simple insertion of a t haplotype-specific element into protein-coding exon II. *Genomics* **28**: 212-219
- Maecker H.T., Todd S.C., Levy S. (1997). The tetraspanin superfamily: molecular facilitators. *Faseb J* **11**: 428-442
- Maekawa M., Sugano K., Kashiwabara H., Ushijima M., Fujita S., Yoshimori M., Kakizoe T. (1999). DNA methylation analysis using bisulfite treatment and PCR-single-strand conformation polymorphism in colorectal cancer showing microsatellite instability. *Biochem Biophys Res Commun* **262**: 671-676
- Maekawa M., Sugano K., Ushijima M., Fukayama N., Nomoto K., Kashiwabara H., Fujita S., Kakizoe T. (2001). Heterogeneity of DNA methylation status

analyzed by bisulfite-PCR-SSCP and correlation with clinico-pathological characteristics in colorectal cancer. *Clin Chem Lab Med* **39**: 121-128

Mai M., Qian C., Yokomizo A., Tindall D.J., Bostwick D., Polychronakos C., Smith D.I., Liu W. (1998). Loss of imprinting and allele switching of p73 in renal cell carcinoma. *Oncogene* **17**: 1739-1741

Mannion B.A., Berditchevski F., Kraeft S.K., Chen L.B., Hemler M.E. (1996). Transmembrane-4 superfamily proteins CD81 (TAPA-1), CD82, CD63, and CD53 specifically associated with integrin alpha 4 beta 1 (CD49d/CD29). *J Immunol* **157**: 2039-2047

Marken J.S., Bajorath J., Edwards C.P., Farr A.G., Schieven G.L., Hellstrom I., Hellstrom K.E., Aruffo A. (1994). Membrane topology of the L6 antigen and identification of the protein epitope recognized by the L6 monoclonal antibody. *J Biol Chem* **269**: 7397-7401

Marra M., Hillier L., Allen M., Bowles M., Dietrich N., Dubuque T., Geisel S., Kucaba T., Lacy M., Le M., Martin J., Morris M., Schellenberg K., Steptoe M., Tan F., Underwood K., Moore B., Theising B., Wylie T., Lennon G., Soares B., Wilson R., Waterston R. (1996). The WashU-HHMI Mouse EST Project. *Unpublished*

Marreiros A., Czolij R., Yardley G., Crossley M., Jackson P. (2003). Identification of regulatory regions within the KAI1 promoter: a role for binding of AP1, AP2 and p53. *Gene* **302**: 155-164

Martin-Alonso J.M., Hernando N., Ghosh S., Coca-Prados M. (1992). Molecular cloning of the bovine CD9 antigen from ocular ciliary epithelial cells. *J Biochem (Tokyo)* **112**: 63-67

Mashimo T., Bandyopadhyay S., Goodarzi G., Watabe M., Pai S.K., Gross S.C., Watabe K. (2000). Activation of the tumor metastasis suppressor gene, KAI1, by etoposide is mediated by p53 and c-Jun genes. *Biochem Biophys Res Commun* **274**: 370-376

Mashimo T., Watabe M., Hirota S., Hosobe S., Miura K., Tegtmeyer P.J., Rinker-Shaeffer C.W., Watabe K. (1998). The expression of the KAI1 gene, a tumor metastasis suppressor, is directly activated by p53. *Proc Natl Acad Sci U S A* **95**: 11307-11311

Mason T.J., Walsh W.P., Lee K., Vogler W. (1992). New opportunities for screening and early detection of bladder cancer. *J Cell Biochem Suppl* **16I**: 13-22

Mau B., Newton M.A., Larget B. (1997). Bayesian phylogenetic inference via Markov chain Monte Carlo methods. *Molecular Biology and Evolution* **14**: 717-724

McCredie M. (1994). Bladder and kidney cancers. *Cancer Surv* **19-20**: 343-368

Meltzer P.S., Jankowski S.A., Dal Cin P., Sandberg A.A., Paz I.B., Coccia M.A. (1991). Identification and cloning of a novel amplified DNA sequence in

human malignant fibrous histiocytoma derived from a region of chromosome 12 frequently rearranged in soft tissue tumors. *Cell Growth Differ* **2**: 495-501

Meola A., Sbardellati A., Bruni Ercole B., Cerretani M., Pezzanera M., Ceccacci A., Vitelli A., Levy S., Nicosia A., Traboni C., McKeating J., Scarselli E. (2000). Binding of hepatitis C virus E2 glycoprotein to CD81 does not correlate with species permissiveness to infection. *J Virol* **74**: 5933-5938

Merika M., Orkin S.H. (1993). DNA-binding specificity of GATA family transcription factors. *Mol Cell Biol* **13**: 3999-4010

Merlo A., Herman J.G., Mao L., Lee D.J., Gabrielson E., Burger P.C., Baylin S.B., Sidransky D. (1995). 5' CpG island methylation is associated with transcriptional silencing of the tumour suppressor p16/CDKN2/MTS1 in human cancers. *Nat Med* **1**: 686-692

Metzelaar M.J., Wijngaard P.L., Peters P.J., Sixma J.J., Nieuwenhuis H.K., Clevers H.C. (1991). CD63 antigen. A novel lysosomal membrane glycoprotein, cloned by a screening procedure for intracellular antigens in eukaryotic cells. *J Biol Chem* **266**: 3239-3245

Miller C., Gurd J., Brass A. (1999). A RAPID algorithm for sequence database comparisons: application to the identification of vector contamination in the EMBL databases. *Bioinformatics* **15**: 111-121

Miller S.A., Dykes D.D., Polesky H.F. (1988). A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* **16**: 1215

Minsky B.D., Chlapowski F.J. (1978). Morphometric analysis of the translocation of luminal membrane between cytoplasm and cell surface of transitional epithelial cells during the expansion-contraction cycles of mammalian urinary bladder. *J Cell Biol* **77**: 685-697

Mitani K. (2002). Molecular mechanisms in leukemogenesis. *Gan To Kagaku Ryoho* **29**: 1107-1112

Miyake M., Koyama M., Seno M., Ikeyama S. (1991). Identification of the motility-related protein (MRP-1), recognized by monoclonal antibody M31-15, which inhibits cell motility. *J Exp Med* **174**: 1347-1354

Miyake M., Nakano K., Ieki Y., Adachi M., Huang C.L., Itoi S., Koh T., Taki T. (1995). Motility related protein 1 (MRP-1/CD9) expression: inverse correlation with metastases in breast cancer. *Cancer Res* **55**: 4127-4131

Miyake M., Nakano K., Itoi S.I., Koh T., Taki T. (1996). Motility-related protein-1 (MRP-1/CD9) reduction as a factor of poor prognosis in breast cancer. *Cancer Res* **56**: 1244-1249

Miyamoto H., Homma M., Hotta H. (1994). Molecular cloning of the murine homologue of CD63/ME491 and detection of its strong expression in the kidney and activated macrophages. *Biochim Biophys Acta* **1217**: 312-316

- Miyamoto H., Kubota Y., Fujinami K., Dobashi Y., Kondo K., Yao M., Shuin T., Hosaka M. (1995). Infrequent somatic mutations of the p16 and p15 genes in human bladder cancer: p16 mutations occur only in low-grade and superficial bladder cancers. *Oncol Res* **7**: 327-330
- Miyamoto H., Kubota Y., Shuin T., Torigoe S., Hosaka M., Iwasaki Y., Danenberg K., Danenberg P.V. (1993). Analyses of p53 gene mutations in primary human bladder cancer. *Oncol Res* **5**: 245-249
- Miyamoto H., Shuin T., Ikeda I., Hosaka M., Kubota Y. (1996). Loss of heterozygosity at the p53, RB, DCC and APC tumor suppressor gene loci in human bladder cancer. *J Urol* **155**: 1444-1447
- Modugno M., Tagliabue E., Ardini E., Berno V., Galmozzi E., De Bortoli M., Castronovo V., Menard S. (2002). p53-dependent downregulation of metastasis-associated laminin receptor. *Oncogene* **21**: 7478-7487
- Moghrabi W.N., Kedzierski W., Travis G.H. (1995). Canine homolog and exclusion of retinal degeneration slow (rds) as the gene for early retinal degeneration (erd) in the dog. *Exp Eye Res* **61**: 641-643
- Moll R., Wu X.R., Lin J.H., Sun T.T. (1995). Uroplakins, specific membrane proteins of urothelial umbrella cells, as histological markers of metastatic transitional cell carcinomas. *Am J Pathol* **147**: 1383-1397
- Momparler R.L., Ayoub J. (2001). Potential of 5-aza-2'-deoxycytidine (Decitabine) a potent inhibitor of DNA methylation for therapy of advanced non-small cell lung cancer. *Lung Cancer* **34**: S111-115
- Momparler R.L., Goodman J. (1977). In vitro cytotoxic and biochemical effects of 5-aza-2'-deoxycytidine. *Cancer Res* **37**: 1636-1639
- Monaco A.P., Larin Z. (1994). YACs, BACs, PACs and MACs: artificial chromosomes as research tools. *Trends Biotechnol* **12**: 280-286
- Morison I.M., Reeve A.E. (1998). A catalogue of imprinted genes and parent-of-origin effects in humans and animals. *Hum Mol Genet* **7**: 1599-1609
- Moses H.L., Pietenpol J.A., Munger K., Murphy C.S., Yang E.Y. (1991). TGF beta regulation of epithelial cell proliferation: role of tumor suppressor genes. *Princess Takamatsu Symp* **22**: 183-195
- Mount S.M. (1982). A catalogue of splice junction sequences. *Nucleic Acids Res* **10**: 459-472
- Muller-Ladner U., Gay R.E., Gay S. (2002). Role of nuclear factor kappaB in synovial inflammation. *Curr Rheumatol Rep* **4**: 201-207
- Muller-Pillasch F., Wallrapp C., Lacher U., Friess H., Buchler M., Adler G., Gress T.M. (1998). Identification of a new tumour-associated antigen TM4SF5 and its expression in human cancer. *Gene* **208**: 25-30 <http://www.elsevier.com>:

80/cgi-bin/cas/tree/store/gene/cas_sub/browse/browse.cgi?year=1998&volume=208&issue=1&aid=11099

- Muneyuki T., Watanabe M., Yamanaka M., Shiraishi T., Isaji S. (2001). KAI1/CD82 expression as a prognostic factor in sporadic colorectal cancer. *Anticancer Res* **21**: 3581-3587
- Murakami T., Li X., Gong J., Bhatia U., Traganos F., Darzynkiewicz Z. (1995). Induction of apoptosis by 5-azacytidine: drug concentration-dependent differences in cell cycle specificity. *Cancer Res* **55**: 3093-3098
- Nagira M., Imai T., Ishikawa I., Uwabe K.I., Yoshie O. (1994). Mouse homologue of C33 antigen (CD82), a member of the transmembrane 4 superfamily: complementary DNA, genomic structure, and expression. *Cell Immunol* **157**: 144-157
- Nan X., Campoy F.J., Bird A. (1997). MeCP2 is a transcriptional repressor with abundant binding sites in genomic chromatin. *Cell* **88**: 471-481
- Nan X., Cross S., Bird A. (1998a). Gene silencing by methyl-CpG-binding proteins. *Novartis Found Symp* **214**: 6-16; discussion 16-21, 46-50
- Nan X., Ng H.H., Johnson C.A., Laherty C.D., Turner B.M., Eisenman R.N., Bird A. (1998b). Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* **393**: 386-389
- Nesci E., Tessitore V. (1969). Scanning electronic microscopic study of the transitional epithelium of the urinary bladder. *Biol Lat* **22**: 235-255
- Ng H.H., Bird A. (1999). DNA methylation and chromatin modification. *Curr Opin Genet Dev* **9**: 158-163
- Nicholas K.B., Nicholas H.B. (2000). Genedoc v2.6.002. <http://www.psc.edu/biomed/genedoc>
- Nishibori M., Cham B., McNicol A., Shalev A., Jain N., Gerrard J.M. (1993). The protein CD63 is in platelet dense granules, is deficient in a patient with Hermansky-Pudlak syndrome, and appears identical to granulophysin. *J Clin Invest* **91**: 1775-1782
- Nishikata H., Oliver C., Mergenhagen S.E., Siraganian R.P. (1992). The rat mast cell antigen AD1 (homologue to human CD63 or melanoma antigen ME491) is expressed in other cells in culture. *J Immunol* **149**: 862-870
- Nishiyama H., Takahashi T., Kakehi Y., Habuchi T., Knowles M.A. (1999). Homozygous deletion at the 9q32-33 candidate tumor suppressor locus in primary human bladder cancer. *Genes Chromosomes Cancer* **26**: 171-175
- Nomoto K., Maekawa M., Sugano K., Ushijima M., Fukayama N., Fujita S., Kakizoe T. (2002). Methylation status and expression of human telomerase reverse transcriptase mRNA in relation to hypermethylation of the p16 gene in

colorectal cancers as analyzed by bisulfite PCR-SSCP. *Jpn J Clin Oncol* **32**: 3-8

Nye J.A., Petersen J.M., Gunther C.V., Jonsen M.D., Graves B.J. (1992). Interaction of murine ets-1 with GGA-binding sites establishes the ETS domain as a new DNA-binding motif. *Genes Dev* **6**: 975-990

Okajima E., Fukuda T., Okita S., Tsutsumi M., Hirao Y., Konishi Y. (1996). Infrequent somatic alteration of p16/MTS1 in human primary superficial bladder cancers. *Cancer Lett* **103**: 227-231

Okano M., Xie S., Li E. (1998). Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. *Nat Genet* **19**: 219-220

Okochi H., Kato M., Nashiro K., Yoshie O., Miyazono K., Furue M. (1997). Expression of tetra-spans transmembrane family (CD9, CD37, CD53, CD63, CD81 and CD82) in normal and neoplastic human keratinocytes: an association of CD9 with alpha 3 beta 1 integrin. *Br J Dermatol* **137**: 856-863

Olsburgh J., Harnden P., Weeks R., Smith B., Joyce A., Hall G., Poulsom R., Selby P., Southgate J. (2003). Uroplakin gene expression in normal human tissues and locally advanced bladder cancer. *J Pathol* **199**: 41-49

Olsburgh J., Weeks R., Selby P., Southgate J. (2002). Human uroplakin lb gene structure and promoter analysis. *Biochim Biophys Acta* **1576**: 163-170

Oren R., Takahashi S., Doss C., Levy R., Levy S. (1990). TAPA-1, the target of an antiproliferative antibody, defines a new family of transmembrane proteins. *Mol Cell Biol* **10**: 4007-4015

Orlow I., Lacombe L., Hannon G.J., Serrano M., Pellicer I., Dalbagni G., Reuter V.E., Zhang Z.F., Beach D., Cordon-Cardo C. (1995). Deletion of the p16 and p15 genes in human bladder tumors. *J Natl Cancer Inst* **87**: 1524-1529

Orlow I., Lianes P., Lacombe L., Dalbagni G., Reuter V.E., Cordon-Cardo C. (1994). Chromosome 9 allelic losses and microsatellite alterations in human bladder tumors. *Cancer Res* **54**: 2848-2851

Ow K., Delprado W., Fisher R., Barrett J., Yu Y., Jackson P., Russell P.J. (2000). Relationship between expression of the KAI1 metastasis suppressor and other markers of advanced bladder cancer. *J Pathol* **191**: 39-47

Oyasu R., Nan L., Szumel R.C., Kawamata H., Hirohashi S. (1995). p53 gene mutations in human urothelial carcinomas: analysis by immunohistochemistry and single-strand conformation polymorphism. *Mod Pathol* **8**: 170-176

Page R.D.M. (1998). TreeView v1.5.0, win32 University of Glasgow, Glasgow, UK. <http://taxonomy.zoology.gla.ac.uk/rod/rod.html>

Page-McCaw A.W., Tsang G., Rubin G.M. (1999). Sequencing Drosophila cDNAs related to tetraspanins. *GenBank direct submission*

- Pak J., Pu Y., Zhang Z.T., Hasty D.L., Wu X.R. (2001). Tamm-Horsfall protein binds to type 1 fimbriated Escherichia coli and prevents E. coli from binding to uroplakin Ia and Ib receptors. *J Biol Chem* **276**: 9924-9930
- Pardee A.B. (1989). G1 events and regulation of cell proliferation. *Science* **246**: 603-608
- Peng L., Li J.Z., Wu H.Z., Wang M.J. (1997). The activation of human platelets mediated by two monoclonal antibodies raised against CD9. *Thromb Res* **87**: 65-73
- Perron J.C., Bixby J.L. (1999). Tetraspanins expressed in the embryonic chick nervous system. *FEBS Lett* **461**: 86-90
- Pichon B., Mercan D., Pouillon V., Christophe-Hobertus C., Christophe D. (2000). A method for the large-scale cloning of nuclear proteins and nuclear targeting sequences on a functional basis. *Anal Biochem* **284**: 231-239
- Pierce J.C., Sauer B., Sternberg N. (1992). A positive selection vector for cloning high molecular weight DNA by the bacteriophage P1 system: improved cloning efficacy. *Proc Natl Acad Sci U S A* **89**: 2056-2060
- Pileri P., Uematsu Y., Campagnoli S., Galli G., Falugi F., Petracca R., Weiner A.J., Houghton M., Rosa D., Grandi G., Abrignani S. (1998). Binding of hepatitis C virus to CD81. *Science* **282**: 938-941
- Pinto A., Attadia V., Fusco A., Ferrara F., Spada O.A., Di Fiore P.P. (1984). 5-Aza-2'-deoxycytidine induces terminal differentiation of leukemic blasts from patients with acute myeloid leukemias. *Blood* **64**: 922-929
- Pinto A., Zagonel V. (1993). 5-Aza-2'-deoxycytidine (Decitabine) and 5-azacytidine in the treatment of acute myeloid leukemias and myelodysplastic syndromes: past, present and future trends. *Leukemia* **7**: 51-60
- Pradhan S., Talbot D., Sha M., Benner J., Hornstra L., Li E., Jaenisch R., Roberts R.J. (1997). Baculovirus-mediated expression and characterization of the full-length murine DNA methyltransferase. *Nucleic Acids Res* **25**: 4666-4673
- Prendergast G.C., Ziff E.B. (1991). Methylation-sensitive sequence-specific DNA binding by the c-Myc basic region. *Science* **251**: 186-189
- Puls K.L., Ni J., Liu D., Morahan G., Wright M.D. (1999). The molecular characterisation of a novel tetraspanin protein, TM4-B(1). *Biochim Biophys Acta* **1447**: 93-99
- Quandt K., Frech K., Karas H., Wingender E., Werner T. (1995). MatInd and MatInspector: new fast and versatile tools for detection of consensus matches in nucleotide sequence data. *Nucleic Acids Res* **23**: 4878-4884 <http://transfac.gbf.de/cgi-bin/matSearch/matsearch.pl>

- Radford K.J., Mallesch J., Hersey P. (1995). Suppression of human melanoma cell growth and metastasis by the melanoma-associated antigen CD63 (ME491). *Int J Cancer* **62**: 631-635
- Rapp G., Freudenstein J., Klaudiny J., Mucha J., Wempe F., Zimmer M., Scheit K.H. (1990). Characterization of three abundant mRNAs from human ovarian granulosa cells. *DNA Cell Biol* **9**: 479-485
- Reese M.G. (1994). Novel neural network algorithms for improved eukaryotic promoter site recognition. German Cancer Research Center, Heidelberg, Germany. http://www.fruitfly.org/cgi-bin/seq_tools/promoter.pl
- Reese M.G., Eeckman F.H. (1995). Novel neural network algorithms for improved eukaryotic promoter site recognition. In "The Seventh International Genome Sequencing and Analysis Conference"., Hilton Head Island, South Carolina, USA.
- Reese M.G., Harris N.L., Eeckman F.H. (1996). Large scale sequencing specific neural networks for promoter and splice site recognition. In "Biocomputing: Proceedings of the 1996 Pacific Symposium". (Hunter L., Klein T.E., Eds.) World Scientific Publishing Co., Singapore.
- Reid L.H., Davies C., Cooper P.R., Crider-Miller S.J., Sait S.N., Nowak N.J., Evans G., Stanbridge E.J., deJong P., Shows T.B., Weissman B.E., Higgins M.J. (1997). A 1-Mb physical map and PAC contig of the imprinted domain in 11p15.5 that contains TAPA1 and the BWSCR1/WT2 region. *Genomics* **43**: 366-375
- Reik W., Surani A. (1997). Genomic Imprinting. IRL Press at Oxford University Press, Oxford, UK.
- Rew D.A. (2001). DNA microarray technology in cancer research. *Eur J Surg Oncol* **27**: 504-508
- Reynolds S.R., Shoemaker C.B., Harn D.A. (1992). T and B cell epitope mapping of SM23, an integral membrane protein of Schistosoma mansoni. *J Immunol* **149**: 3995-4001
- Reznikoff C.A., Belair C.D., Yeager T.R., Savelieva E., Blelloch R.H., Puthenveettil J.A., Cuthill S. (1996). A molecular genetic model of human bladder cancer pathogenesis. *Semin Oncol* **23**: 571-584
- Rhee I., Bachman K.E., Park B.H., Jair K.W., Yen R.W., Schuebel K.E., Cui H., Feinberg A.P., Lengauer C., Kinzler K.W., Baylin S.B., Vogelstein B. (2002). DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. *Nature* **416**: 552-556
- Roberts A.B. (2002). The ever-increasing complexity of TGF-beta signaling. *Cytokine Growth Factor Rev* **13**: 3-5
- Robertson K.D., Uzvolgyi E., Liang G., Talmadge C., Sumegi J., Gonzales F.A., Jones P.A. (1999). The human DNA methyltransferases (DNMTs) 1, 3a and

- 3b: coordinate mRNA expression in normal tissues and overexpression in tumors. *Nucleic Acids Res* **27**: 2291-2298
- Rollins B.J., Stiles C.D. (1989). Serum-inducible genes. *Adv Cancer Res* **53**: 1-32
- Rosin M.P., Anwar W. (1992). Chromosomal damage in urothelial cells from Egyptians with chronic Schistosoma haematobium infections. *Int J Cancer* **50**: 539-543
- Ross A.H., Dietzschold B., Jackson D.M., Earley J.J., Jr., Ghrist B.D., Atkinson B., Koprowski H. (1985). Isolation and amino terminal sequencing of a novel melanoma-associated antigen. *Arch Biochem Biophys* **242**: 540-548
- Roth S.Y., Denu J.M., Allis C.D. (2001). Histone acetyltransferases. *Annu Rev Biochem* **70**: 81-120
- Rubinstein E., Billard M., Plaisance S., Prenant M., Boucheix C. (1993). Molecular cloning of the mouse equivalent of CD9 antigen. *Thromb Res* **71**: 377-383
- Rubinstein E., Le Naour F., Billard M., Prenant M., Boucheix C. (1994). CD9 antigen is an accessory subunit of the VLA integrin complexes. *Eur J Immunol* **24**: 3005-3013
- Rubinstein E., Le Naour F., Lagaudriere-Gesbert C., Billard M., Conjeaud H., Boucheix C. (1996). CD9, CD63, CD81, and CD82 are components of a surface tetraspan network connected to HLA-DR and VLA integrins. *Eur J Immunol* **26**: 2657-2665
- Saffer J.D., Jackson S.P., Annarella M.B. (1991). Developmental expression of Sp1 in the mouse. *Mol Cell Biol* **11**: 2189-2199
- Salem C., Liang G., Tsai Y.C., Coulter J., Knowles M.A., Feng A.C., Groshen S., Nichols P.W., Jones P.A. (2000). Progressive increases in de novo methylation of CpG islands in bladder cancer. *Cancer Res* **60**: 2473-2476
- Sambrook J., Fritsch E.F., Maniatis T. (1989). Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA.
- Sandal T. (2002). Molecular aspects of the mammalian cell cycle and cancer. *Oncologist* **7**: 73-81
- Sarikas S.N., Chlapowski F.J. (1986). Effect of ATP inhibitors on the translocation of luminal membrane between cytoplasm and cell surface of transitional epithelial cells during the expansion-contraction cycle of the rat urinary bladder. *Cell Tissue Res* **246**: 109-117
- Sarikas S.N., Chlapowski F.J. (1989). The effect of thioglycolate on intermediate filaments and membrane translocation in rat urothelium during the expansion-contraction cycle. *Cell Tissue Res* **258**: 393-401
- Schmidt C., Kunemund V., Wintergerst E.S., Schmitz B., Schachner M. (1996). CD9 of mouse brain is implicated in neurite outgrowth and cell migration in vitro and

is associated with the alpha 6/beta 1 integrin and the neural adhesion molecule L1. *J Neurosci Res* **43**: 12-31

Schmidt W.M., Mueller M.W. (1999). CapSelect: a highly sensitive method for 5' CAP-dependent enrichment of full-length cDNA in PCR-mediated analysis of mRNAs. *Nucleic Acids Res* **27**: e31

Schug J., Overton G.C. (1997). TESS: Transcription element search software on the WWW; Technical Report CBIL-TR-1997-1001-v0.0, of the Computational Biology and Informatics Laboratory. School of Medicine, University of Pennsylvania, Philadelphia, PA, USA. <http://www.cbil.upenn.edu/tess>

Sekita N., Suzuki H., Ichikawa T., Kito H., Akakura K., Igarashi T., Nakayama T., Watanabe M., Shiraishi T., Toyota M., Yoshie O., Ito H. (2001). Epigenetic regulation of the KAI1 metastasis suppressor gene in human prostate cancer cell lines. *Jpn J Cancer Res* **92**: 947-951

Selvey S., Thompson E.W., Matthaei K., Lea R.A., Irving M.G., Griffiths L.R. (2001). Beta-actin: an unsuitable internal control for RT-PCR. *Mol Cell Probes* **15**: 307-311

Seraj M.J., Thomas A.R., Chin J.L., Theodorescu D. (2001). Molecular determination of perivesical and lymph node metastasis after radical cystectomy for urothelial carcinoma of the bladder. *Clin Cancer Res* **7**: 1516-1522

Serazin-Leroy V., Denis-Henriot D., Morot M., de Mazancourt P., Giudicelli Y. (1998). Semi-quantitative RT-PCR for comparison of mRNAs in cells with different amounts of housekeeping gene transcripts. *Mol Cell Probes* **12**: 283-291

Serru V., Dessen P., Boucheix C., Rubinstein E. (2000). Sequence and expression of seven new tetraspans. *Biochim Biophys Acta* **1478**: 159-163

Serru V., Le Naour F., Billard M., Azorsa D.O., Lanza F., Boucheix C., Rubinstein E. (1999). Selective tetraspan-integrin complexes (CD81/alpha4beta1, CD151/alpha3beta1, CD151/alpha6beta1) under conditions disrupting tetraspan interactions. *Biochem J* **340**: 103-111

Shanmukhappa K., Kapil S. (2000). CD151/PETA-3, a tetraspanin molecule, interacts with the 3' untranslated region and partial nucleoprotein gene of Porcine Reproductive and Respiratory Syndrome Virus RNA. *GenBank direct submission*

Sho M., Adachi M., Taki T., Hashida H., Konishi T., Huang C.L., Ikeda N., Nakajima Y., Kanehiro H., Hisanaga M., Nakano H., Miyake M. (1998). Transmembrane 4 superfamily as a prognostic factor in pancreatic cancer. *Int J Cancer* **79**: 509-516

Sidransky D., Frost P., Von Eschenbach A., Oyasu R., Preisinger A.C., Vogelstein B. (1992). Clonal origin bladder cancer. *N Engl J Med* **326**: 737-740

- Simon D., Larget B. (2001). Bayesian analysis in molecular biology and evolution v2.03 Duquesne University, Pittsburgh, PA, USA. <http://www.mathcs.duq.edu/lerget/bambe.html>
- Sincock P.M., Mayrhofer G., Ashman L.K. (1997). Localization of the transmembrane 4 superfamily (TM4SF) member PETA-3 (CD151) in normal human tissues: comparison with CD9, CD63, and alpha5beta1 integrin. *J Histochem Cytochem* **45**: 515-525
- Singal R., Ginder G.D. (1999). DNA methylation. *Blood* **93**: 4059-4070
- Singer J., Roberts-Ems J., Riggs A.D. (1979). Methylation of mouse liver DNA studied by means of the restriction enzymes msp I and hpa II. *Science* **203**: 1019-1021
- Smith H.O., Birnstiel M.L. (1976). A simple method for DNA restriction site mapping. *Nucleic Acids Res* **3**: 2387-2398
- Sohma Y., Suzuki T., Sasano H., Nagura H., Nose M., Yamamoto T. (1994). Increased mRNA for CD63 antigen in atherosclerotic lesions of Watanabe heritable hyperlipidemic rabbits. *Cell Struct Funct* **19**: 219-225
- Sorlie T., Martel-Planche G., Hainaut P., Lewalter J., Holm R., Borresen-Dale A.L., Montesano R. (1998). Analysis of p53, p16MTS, p21WAF1 and H-ras in archived bladder tumours from workers exposed to aromatic amines. *Br J Cancer* **77**: 1573-1579
- Soulitzis N., Sourvinos G., Dokianakis D.N., Spandidos D.A. (2002). p53 codon 72 polymorphism and its association with bladder cancer. *Cancer Lett* **179**: 175-183
- South A.C.R. (2001). 2001 Cancer Report: Epidemiology of Cancer in South Australia 1977-2000, Department of Human Services, Adelaide, Australia
- Southern E.M. (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* **98**: 503-517
- Spruck C.H., 3rd, Ohneseit P.F., Gonzalez-Zulueta M., Esrig D., Miyao N., Tsai Y.C., Lerner S.P., Schmutte C., Yang A.S., Cote R. (1994). Two molecular pathways to transitional cell carcinoma of the bladder. *Cancer Res* **54**: 784-788
- Spruck C.H., 3rd, Rideout W.M., 3rd, Olumi A.F., Ohneseit P.F., Yang A.S., Tsai Y.C., Nichols P.W., Horn T., Hermann G.G., Steven K. (1993). Distinct pattern of p53 mutations in bladder cancer: relationship to tobacco usage. *Cancer Res* **53**: 1162-1166
- Stadler W.M., Olopade O.I. (1996). The 9p21 region in bladder cancer cell lines: large homozygous deletion inactivate the CDKN2, CDKN2B and MTAP genes. *Urol Res* **24**: 239-244

- Staehelin L.A., Chlapowski F.J., Bonneville M.A. (1972). Luminal plasma membrane of the urinary bladder. I. Three-dimensional reconstruction from freeze-etch images. *J Cell Biol* **53**: 73-91
- Stegmaier K., Pendse S., Barker G.F., Bray-Ward P., Ward D.C., Montgomery K.T., Krauter K.S., Reynolds C., Sklar J., Donnelly M. (1995). Frequent loss of heterozygosity at the TEL gene locus in acute lymphoblastic leukemia of childhood. *Blood* **86**: 38-44
- Sullivan C.D., Geisert E.E., Jr. (1998). Expression of rat target of the antiproliferative antibody (TAPA) in the developing brain. *J Comp Neurol* **396**: 366-380
- Sun T.T., Zhao H., Provet J., Aebi U., Wu X.R. (1996). Formation of asymmetric unit membrane during urothelial differentiation. *Mol Biol Rep* **23**: 3-11
- Szala S., Kasai Y., Steplewski Z., Rodeck U., Koprowski H., Linnenbach A.J. (1990). Molecular cloning of cDNA for the human tumor-associated antigen CO-029 and identification of related transmembrane antigens. *Proc Natl Acad Sci U S A* **87**: 6833-6837
- Tachibana I., Bodorova J., Berditchevski F., Zutter M.M., Hemler M.E. (1997). NAG-2, a novel transmembrane-4 superfamily (TM4SF) protein that complexes with integrins and other TM4SF proteins. *J Biol Chem* **272**: 29181-29189
- Tachibana I., Hemler M.E. (1999). Role of transmembrane 4 superfamily (TM4SF) proteins CD9 and CD81 in muscle cell fusion and myotube maintenance. *J Cell Biol* **146**: 893-904
- Tai X.G., Yashiro Y., Abe R., Toyooka K., Wood C.R., Morris J., Long A., Ono S., Kobayashi M., Hamaoka T., Neben S., Fujiwara H. (1996). A role for CD9 molecules in T cell activation. *J Exp Med* **184**: 753-758
- Takagi S., Fujikawa K., Imai T., Fukuhara N., Fukudome K., Minegishi M., Tsuchiya S., Konno T., Hinuma Y., Yoshie O. (1995). Identification of a highly specific surface marker of T-cell acute lymphoblastic leukemia and neuroblastoma as a new member of the transmembrane 4 superfamily. *Int J Cancer* **61**: 706-715
- Takahashi S., Doss C., Levy S., Levy R. (1990). TAPA-1, the target of an antiproliferative antibody, is associated on the cell surface with the Leu-13 antigen. *J Immunol* **145**: 2207-2213
- Takasuga A., Hirotsume S., Itoh R., Jitohzono A., Suzuki H., Aso H., Sugimoto Y. (2001). Establishment of a high throughput EST sequencing system using poly(A) tail-removed cDNA libraries and determination of 36,000 bovine ESTs. *Nucleic Acids Res* **29**: E108
- Taylor P.L. (1990). GeneJockey sequence processor. Biosoft, Cambridge, UK.
- Thiesen H.J., Bach C. (1990). Target Detection Assay (TDA): a versatile procedure to determine DNA binding sites as demonstrated on SP1 protein. *Nucleic Acids Res* **18**: 3203-3209

- Thompson J.D., Higgins D.G., Gibson T.J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* **22**: 4673-4680
- Todd S.C., Doctor V.S., Levy S. (1998). Sequences and expression of six new members of the tetraspanin/TM4SF family. *Biochim Biophys Acta* **1399**: 101-104 http://www.elsevier.com:80/cgi-bin/cas/store/bbaexp/cas_sub/browse/browse.cgi?year=1998&volume=1399&issue=1&aid=91151
- Todd S.C., Lipps S.G., Crisa L., Salomon D.R., Tsoukas C.D. (1996). CD81 expressed on human thymocytes mediates integrin activation and interleukin 2-dependent proliferation. *J Exp Med* **184**: 2055-2060
- Todres E., Nardi J.B., Robertson H.M. (2000). The tetraspanin superfamily in insects. *Insect Mol Biol* **9**: 581-590
- Tomlinson M.G., Wright M.D. (1996). Characterisation of mouse CD37: cDNA and genomic cloning. *Mol Immunol* **33**: 867-872
- Toruner G.A., Ucar A., Tez M., Cetinkaya M., Ozen H., Ozcelik T. (2001). p53 codon 72 polymorphism in bladder cancer--no evidence of association with increased risk or invasiveness. *Urol Res* **29**: 393-395
- Toyota M., Ahuja N., Ohe-Toyota M., Herman J.G., Baylin S.B., Issa J.P. (1999). CpG island methylator phenotype in colorectal cancer. *Proc Natl Acad Sci U S A* **96**: 8681-8686
- Toyota M., Issa J.P. (1999). CpG island methylator phenotypes in aging and cancer. *Semin Cancer Biol* **9**: 349-357
- Travis G.H., Brennan M.B., Danielson P.E., Kozak C.A., Sutcliffe J.G. (1989). Identification of a photoreceptor-specific mRNA encoded by the gene responsible for retinal degeneration slow (rds). *Nature* **338**: 70-73
- Travis G.H., Christerson L., Danielson P.E., Klisak I., Sparkes R.S., Hahn L.B., Dryja T.P., Sutcliffe J.G. (1991). The human retinal degeneration slow (RDS) gene: chromosome assignment and structure of the mRNA. *Genomics* **10**: 733-739
- Tseng C.T., Klimpel G.R. (2002). Binding of the hepatitis C virus envelope protein E2 to CD81 inhibits natural killer cell functions. *J Exp Med* **195**: 43-49
- Tseng C.T., Miskovsky E., Klimpel G.R. (2001). Crosslinking CD81 results in activation of TCRgammadelta T cells. *Cell Immunol* **207**: 19-27
- Tsutsumi M., Sugano K., Yamaguchi K., Kakizoe T., Akaza H. (1997). Correlation of allelic loss of the p53 gene and tumor grade, stage, and malignant progression in bladder cancer. *Int J Urol* **4**: 74-78
- Uzawa K., Ono K., Suzuki H., Tanaka C., Yakushiji T., Yamamoto N., Yokoe H., Tanzawa H. (2002). High prevalence of decreased expression of KAI1

metastasis suppressor in human oral carcinogenesis. *Clin Cancer Res* **8**: 828-835

Van Cong N., Benoit P., Gross M.S., Uzan G., Frachet P., Marguerie G., Frezal J., Boucheix C. (1989). Assignment of the gene for CD9 (p24) antigen to 12p13. *Cytogenet Cell Genet* **51**: 1096

Virtaneva K.I., Angelisova P., Baumruker T., Horejsi V., Nevanlinna H., Schroder J. (1993). The genes for CD37, CD53, and R2, all members of a novel gene family, are located on different chromosomes. *Immunogenetics* **37**: 461-465

Virtaneva K.I., Emi N., Marken J.S., Aruffo A., Jones C., Spurr N.K., Schroder J.P. (1994). Chromosomal localization of three human genes coding for A15, L6, and S5.7 (TAPA1): all members of the transmembrane 4 superfamily of proteins. *Immunogenetics* **39**: 329-334

Wada H., Murakami A., Ito J., Okisaka S. (1996). Molecular cloning and characterization of bovine rom1 cDNA and gene. *GenBank direct submission*

Wada S., Yoshimura R., Masuda C., Hase T., Ikemoto S., Kishimoto T., Fukushima S. (2001). Are tobacco use and urine pH indicated as risk factors for bladder carcinoma? *Int J Urol* **8**: 106-109

Wada T., Louhelainen J., Hemminki K., Adolfsson J., Wijkstrom H., Norming U., Borgstrom E., Hansson J., Sandstedt B., Steineck G. (2000). Bladder cancer: allelic deletions at and around the retinoblastoma tumor suppressor gene in relation to stage and grade. *Clin Cancer Res* **6**: 610-615

Wallace R.B., Shaffer J., Murphy R.F., Bonner J., Hirose T., Itakura K. (1979). Hybridization of synthetic oligodeoxyribonucleotides to phi chi 174 DNA: the effect of single base pair mismatch. *Nucleic Acids Res* **6**: 3543-3557

Wang M.X., Earley J.J., Jr., Shields J.A., Donoso L.A. (1992). An ocular melanoma-associated antigen. Molecular characterization. *Arch Ophthalmol* **110**: 399-404

Wang Q., Chen Q., Zhao K., Wang L., Traboulsi E.I. (2001). Update on the molecular genetics of retinitis pigmentosa. *Ophthalmic Genet* **22**: 133-154

Waterston R. (1998). Genome sequence of the nematode *C. elegans*: a platform for investigating biology. The *C. elegans* Sequencing Consortium. *Science* **282**: 2012-2018

Webb G.C., Finch J.L., Cowled P.A. (1999). Assignment of the Uroplakin 1b (Upk1b) gene to mouse chromosome 16 bands B5-C2 by in situ hybridization. *Cytogenet Cell Genet* **84**: 37-38

Weinel C., Tummler B., Hilbert H., Nelson K.E., Kiewitz C. (2001). General method of rapid Smith/Birnstiel mapping adds for gap closure in shotgun microbial genome sequencing projects: application to *Pseudomonas putida* KT2440. *Nucleic Acids Res* **29**: E110

- Weng J., Belecky-Adams T., Adler R., Travis G.H. (1998). Identification of two rds/peripherin homologs in the chick retina. *Invest Ophthalmol Vis Sci* **39**: 440-443
- Werner H., Dimou L., Klugmann M., Pfeiffer S., Nave K.A. (2001). Multiple splice isoforms of proteolipid M6B in neurons and oligodendrocytes. *Mol Cell Neurosci* **18**: 593-605
- Wessing A., Eichelberg D. (1969). Electron microscopic studies of the renal tubules (malpighian tubules) in *Drosophila melanogaster*. I. Regional formation of tubules. *Z Zellforsch Mikrosk Anat* **101**: 285-322
- Wessing A., Eichelberg D. (1972a). Electron microscopic studies of the renal tubules (Malpighian tubules) in *Drosophila melanogaster*. III. Intracellular storage of the amino-acid 3-hydroxykynurenin. *Z Zellforsch Mikrosk Anat* **125**: 132-142
- Wessing A., Eichelberg D. (1972b). Electron microscopy studies of the renal tubules (malpighian tubules) in *Drosophila melanogaster*. IV. Localization of sodium and chloride ions. *Z Zellforsch Mikrosk Anat* **131**: 269-286
- White A., Lamb P.W., Barrett J.C. (1998). Frequent downregulation of the KAI1(CD82) metastasis suppressor protein in human cancer cell lines. *Oncogene* **16**: 3143-3149
- White O., Dunning T., Sutton G., Adams M., Venter J.C., Fields C. (1993). A quality control algorithm for DNA sequencing projects. *Nucleic Acids Res* **21**: 3829-3838
- Wice B.M., Gordon J.I. (1995). A tetraspan membrane glycoprotein produced in the human intestinal epithelium and liver that can regulate cell density-dependent proliferation. *J Biol Chem* **270**: 21907-21918
- Wilder R.L., Griffiths M.M., Remmers E.F., Cannon G.W., Caspi R.R., Kawahito Y., Gulko P.S., Longman R.E., Dracheva S.V., Du Y., Sun S.H., Wang J., Shepard J.S., Joe B., Ge L., Chen S., Chang L., Hoffman J., Silver P.B., Reese V.R. (1999). Localization in rats of genetic loci regulating susceptibility to experimental erosive arthritis and related autoimmune diseases. *Transplant Proc* **31**: 1585-1588
- Willett B.J., Neil J.C. (1995). cDNA cloning and eukaryotic expression of feline CD9. *Mol Immunol* **32**: 417-423
- Williamson M.P., Elder P.A., Shaw M.E., Devlin J., Knowles M.A. (1995). p16 (CDKN2) is a major deletion target at 9p21 in bladder cancer. *Hum Mol Genet* **4**: 1569-1577
- Wingender E., Chen X., Hehl R., Karas H., Liebich I., Matys V., Meinhardt T., Pruss M., Reuter I., Schacherer F. (2000). TRANSFAC: an integrated system for gene expression regulation. *Nucleic Acids Res* **28**: 316-319
- Wright M.D., Rochelle J.M., Tomlinson M.G., Seldin M.F., Williams A.F. (1993). Gene structure, chromosomal localization, and protein sequence of mouse CD53

(Cd53): evidence that the transmembrane 4 superfamily arose by gene duplication. *Int Immunol* **5**: 209-216

Wright M.D., Tomlinson M.G. (1994). The ins and outs of the transmembrane 4 superfamily. *Immunol Today* **15**: 588-594

Wu R.L., Osman I., Wu X.R., Lu M.L., Zhang Z.F., Liang F.X., Hamza R., Scher H., Cordon-Cardo C., Sun T.T. (1998). Uroplakin II gene is expressed in transitional cell carcinoma but not in bilharzial bladder squamous cell carcinoma: alternative pathways of bladder epithelial differentiation and tumor formation. *Cancer Res* **58**: 1291-1297

Wu W.J., Kakehi Y., Chang S.F., Huang C.H., Chang L.L. (2000). Genetic alterations of the p16 gene in urothelial carcinoma in Taiwanese patients. *BJU Int* **85**: 143-149

Wu X.R., Lin J.H., Walz T., Haner M., Yu J., Aebi U., Sun T.T. (1994). Mammalian uroplakins. A group of highly conserved urothelial differentiation-related membrane proteins. *J Biol Chem* **269**: 13716-13724

Wu X.R., Manabe M., Yu J., Sun T.T. (1990). Large scale purification and immunolocalization of bovine uroplakins I, II, and III. Molecular markers of urothelial differentiation. *J Biol Chem* **265**: 19170-19179

Wu X.R., Medina J.J., Sun T.T. (1995). Selective interactions of UPIa and UPIb, two members of the transmembrane 4 superfamily, with distinct single transmembrane-domained proteins in differentiated urothelial cells. *J Biol Chem* **270**: 29752-29759

Wu X.R., Sun T.T. (1993). Molecular cloning of a 47 kDa tissue-specific and differentiation-dependent urothelial cell surface glycoprotein. *J Cell Sci* **106**: 31-43

Wu X.R., Sun T.T., Medina J.J. (1996). In vitro binding of type 1-fimbriated *Escherichia coli* to uroplakins Ia and Ib: relation to urinary tract infections. *Proc Natl Acad Sci USA* **93**: 9630-9635

Xiong Z., Laird P.W. (1997). COBRA: a sensitive and quantitative DNA methylation assay. *Nucleic Acids Res* **25**: 2532-2534

Xu G.L., Bestor T.H., Bourc'his D., Hsieh C.L., Tommerup N., Bugge M., Hulten M., Qu X., Russo J.J., Viegas-Pequignot E. (1999). Chromosome instability and immunodeficiency syndrome caused by mutations in a DNA methyltransferase gene. *Nature* **402**: 187-191

Yamada H., Chen D., Monstein H.J., Hakanson R. (1997). Effects of fasting on the expression of gastrin, cholecystokinin, and somatostatin genes and of various housekeeping genes in the pancreas and upper digestive tract of rats. *Biochem Biophys Res Commun* **231**: 835-838

Yanez-Mo M., Mittelbrunn M., Sanchez-Madrid F. (2001). Tetraspanins and intercellular interactions. *Microcirculation* **8**: 153-168

- Yen R.W., Vertino P.M., Nelkin B.D., Yu J.J., el-Deiry W., Cumaraswamy A., Lennon G.G., Trask B.J., Celano P., Baylin S.B. (1992). Isolation and characterization of the cDNA encoding human DNA methyltransferase. *Nucleic Acids Res* **20**: 2287-2289
- Yi E.S., Shabaik A.S., Lacey D.L., Bedoya A.A., Yin S., Housley R.M., Danilenko D.M., Benson W., Cohen A.M., Pierce G.F., Thomason A., Ulich T.R. (1995). Keratinocyte growth factor causes proliferation of urothelium in vivo. *J Urol* **154**: 1566-1570
- Yoder J.A., Bestor T.H. (1998). A candidate mammalian DNA methyltransferase related to pmt1p of fission yeast. *Hum Mol Genet* **7**: 279-284
- Yu J., Lin J.H., Wu X.R., Sun T.T. (1994). Uroplakins Ia and Ib, two major differentiation products of bladder epithelium, belong to a family of four transmembrane domain (4TM) proteins. *J Cell Biol* **125**: 171-182
- Yu J., Manabe M., Sun T.T. (1992). Identification of an 85-100 kDa glycoprotein as a cell surface marker for an advanced stage of urothelial differentiation: association with the inter-plaque ('hinge') area. *Epithelial Cell Biol* **1**: 4-12
- Yu J., Manabe M., Wu X.R., Xu C., Surya B., Sun T.T. (1990). Uroplakin I: a 27-kD protein associated with the asymmetric unit membrane of mammalian urothelium. *J Cell Biol* **111**: 1207-1216
- Yu Y., Yang J.L., Markovic B., Jackson P., Yardley G., Barrett J., Russell P.J. (1997). Loss of KAI1 messenger RNA expression in both high-grade and invasive human bladder cancers. *Clin Cancer Res* **3**: 1045-1049
- Yuasa T., Yoshiki T., Isono T., Tanaka T., Hayashida H., Okada Y. (1999). Expression of transitional cell-specific genes, uroplakin Ia and II, in bladder cancer: detection of circulating cancer cells in the peripheral blood of metastatic patients. *Int J Urol* **6**: 286-292
- Yuasa T., Yoshiki T., Tanaka T., Kim C.J., Isono T., Okada Y. (1998). Expression of uroplakin Ib and uroplakin III genes in tissues and peripheral blood of patients with transitional cell carcinoma. *Jpn J Cancer Res* **89**: 879-882
- Zemni R., Bienvenu T., Vinet M.C., Sefiani A., Carrie A., Billuart P., McDonell N., Couvert P., Francis F., Chafey P., Fauchereau F., Friocourt G., Portes V., Cardona A., Frints S., Meindl A., Brandau O., Ronce N., Moraine C., Bokhoven H., Ropers H.H., Sudbrak R., Kahn A., Fryns J.P., Beldjord C., Chelly J. (2000). A new gene involved in X-linked mental retardation identified by analysis of an X;2 balanced translocation. *Nat Genet* **24**: 167-170
- Zhang J., Ramesh N., Chen Y., Li Y., Dilley J., Working P., Yu D.C. (2002). Identification of human uroplakin II promoter and its use in the construction of CG8840, a urothelium-specific adenovirus variant that eliminates established bladder tumors in combination with docetaxel. *Cancer Res* **62**: 3743-3750

Zhang X.Y., Ehrlich K.C., Wang R.Y., Ehrlich M. (1986). Effect of site-specific DNA methylation and mutagenesis on recognition by methylated DNA-binding protein from human placenta. *Nucleic Acids Res* **14**: 8387-8397

Zhou G., Mo W.J., Sebbel P., Min G., Neubert T.A., Glockshuber R., Wu X.R., Sun T.T., Kong X.P. (2001). Uroplakin Ia is the urothelial receptor for uropathogenic Escherichia coli: evidence from in vitro FimH binding. *J Cell Sci* **114**: 4095-410