

09PH  
H9281



**Identification and analysis of genes involved in**  
**cotton fibre initiation**

This thesis is presented for the degree of Doctor of Philosophy  
(PhD) in the Faculty of Sciences at the University of Adelaide

**John A Humphries**

**Discipline of Genetics, School of Molecular and  
Biomedical Science, University of Adelaide**

**September 2006**

## Table of Contents

<b>Declaration</b>	<b>7</b>
<b>Acknowledgements</b>	<b>8</b>
<b>Abstract</b>	<b>9</b>
<b>Chapter 1: Introduction</b>	<b>11</b>
<b>1.1 Cotton crop and industry</b>	<b>11</b>
<b>1.2 Cotton taxonomy</b>	<b>11</b>
<b>1.3 Growth of the cotton plant</b>	<b>12</b>
<b>1.4 Cotton fibre development</b>	<b>12</b>
1.4.1 Fibre initiation	12
1.4.2 Fibre elongation	13
1.4.3 Secondary wall synthesis	13
1.4.4 Maturation	14
<b>1.5 Genes involved in cotton fibre development</b>	<b>14</b>
1.5.1 Genes involved in elongation and secondary cell wall growth	15
1.5.2 Genes involved in initiation of cotton fibres	16
<b>1.6 Cotton Biotechnology</b>	<b>17</b>
1.6.1 Plant breeding and genetic engineering	17
1.6.2 Towards fibre improvement in genetically modified cotton	17
1.6.3 Transformation of cotton	18
<b>1.7 <i>Arabidopsis</i> trichomes</b>	<b>19</b>
<b>1.8 Genetic regulation of <i>Arabidopsis</i> trichome initiation</b>	<b>20</b>
1.8.1 <i>TRANSPARENT TESTA GLABRA1 (TTG1)</i>	21
1.8.2 <i>GLABROUS1 (GL1)</i>	23
1.8.3 <i>GLABRA3 (GL3)</i>	24
1.8.4 <i>TRANSPARENT TESTA8 (TT8)</i> and <i>ENHANCER OF GL3 (EGL3)</i>	24
1.8.5 <i>TRIPTYCHON (TRY)</i> and <i>CAPRICE (CPC)</i>	25
1.8.6 <i>GLABRA2 (GL2)</i>	26

1.8.7 Trichome and root hair initiation	26
1.8.8 The WD40/bHLH/MYB regulatory complex	27
<b>1.9 <i>Arabidopsis</i> trichomes as a model for cotton fibre development</b>	<b>28</b>
<b>1.10 Experimental approach</b>	<b>30</b>
<b>Chapter 2: Materials and Methods</b>	<b>33</b>
<hr/>	
2.1 Plant material	33
2.2 Enzymes	33
2.3 Radioactive isotopes	34
2.4 Molecular weight markers	34
2.5 Antibiotics and indicators	34
2.6 Kits	34
2.8 Chemicals reagents and miscellaneous	35
2.9 Vectors	35
2.10 Yeast and bacterial strains	36
2.11 Oligonucleotides	36
2.12 Solutions, buffers and media	38
2.13 Media	39
2.14 Plant techniques	40
2.15 General molecular techniques	41
2.16 Southern blot analysis	44
2.17 <i>In situ</i> hybridisation	46
2.18 Yeast two-hybrid assay and yeast methods	46
<b>Chapter 3: WD-repeat genes in cotton</b>	<b>49</b>
<hr/>	
3.1 Previous work: Isolation of four WD-repeat genes in cotton with homology to <i>TTG1</i>	49
3.2 Genomic organisation of cotton WD-repeat genes	52

3.2.1 Genomic PCR analysis	52
3.2.2 Southern blot analysis of WD-repeat genes	53
<b>3.3 Phylogenetic analysis of cotton WD-repeat sequences</b>	<b>54</b>
<b>3.4 Discussion</b>	<b>55</b>
3.4.1 Isolation of four <i>TTG1</i> -like genes in cotton	55
3.4.2 All <i>GhTTG</i> genes originate from the ancestral D genome	56
3.4.3 Phylogenetic analysis of WD-repeat genes	57
<b><u>Chapter 4: Cross-species complementation of cotton WD-repeat genes</u></b>	<b>61</b>
<b>4.1 Introduction</b>	<b>61</b>
<b>4.2 Previous work</b>	<b>61</b>
<b>4.3 <i>Arabidopsis</i> stable transformation</b>	<b>62</b>
4.3.1 <i>Agrobacterium</i> -mediated genetic transformation	62
4.3.2 Transformation of <i>Arabidopsis ttg1-1</i> mutant plants	63
4.3.3 Additional phenotypes	65
<b>4.4 Discussion</b>	<b>67</b>
<b>4.5 Summary</b>	<b>71</b>
<b><u>Chapter 5: Temporal and spatial expression of WD-repeat genes in cotton</u></b>	<b>73</b>
<b>5.1 Introduction</b>	<b>73</b>
<b>5.2 Reverse-transcriptase PCR analysis</b>	<b>73</b>
<b>5.3 Cellular localisation of cotton WD-repeat genes within the ovule</b>	<b>74</b>
5.3.1 Technical aspects of <i>in situ</i> hybridisation	74
5.3.2 Generation of <i>in situ</i> hybridisation RNA probes	75
5.3.3 <i>In situ</i> hybridisation results	77
<b>5.4 Analysis of <i>GhTTG</i> expression in fibreless cotton mutants</b>	<b>79</b>
5.4.1 Fibreless mutants of cotton	79
5.4.2 Genomic PCRs on fibreless mutants	80
5.4.3 Reverse transcriptase-PCR analysis of a fibreless mutant	80
5.4.4 Real-time PCR	81

5.5 Discussion	82
<b><u>Chapter 6: Molecular interactions of cotton WD-repeat proteins</u></b>	<b>87</b>
6.1 Introduction	87
6.2 Yeast two-hybrid	87
6.3 Investigation of GhTTG interaction with GL3/EGL3	88
6.4 Screen for interacting proteins of GhTTG in cotton	91
6.5 Interacting Clone #1: NAC domain protein (GhNAC)	95
6.6 Interacting Clone #2: TCP-domain protein (GhTCP)	96
6.7 Discussion	98
6.7.1 WD40 interactions	98
6.7.2 Testing of GhTTG protein interactions	99
6.7.3 A cotton fibre cDNA library screen for interactors of GhTTG3	100
6.8 Summary	102
<b><u>Chapter 7: Discussion and conclusions</u></b>	<b>103</b>
7.1 Summary of results	103
7.2 Further experiments	104
7.3 <i>Arabidopsis</i> trichomes as a model system for cotton fibre development	106
7.4 Implications for cotton fibre development	108
7.5 Conclusion	109
<b><u>Appendix 1: Vector Maps</u></b>	<b>111</b>
<b><u>Appendix 2: Real-time PCR data</u></b>	<b>114</b>
<b><u>Appendix 3: <math>\alpha</math>-galactosidase assay</u></b>	<b>115</b>
<b><u>Appendix 4: Sequence identification numbers</u></b>	<b>116</b>
<b><u>References</u></b>	<b>117</b>

## **Abstract**

Cotton fibres are single-celled hairs, arising from the epidermal surface of the cotton ovule. One factor in determining the length of the mature cotton fibre is the timing of fibre initiation, which is therefore a crucial step in obtaining commercial cotton fibres. To achieve a greater understanding of the regulation of cotton fibre differentiation, more fundamental information is needed on the signals and mechanisms associated with fibre initiation. The extensive genetic knowledge of *Arabidopsis* leaf trichomes could aid in the elucidation of the genetic mechanisms controlling cotton fibre differentiation. Trichomes are small hairs on the plant surface, originating from single epidermal cells in a developmental process that appears very similar to that of cotton fibres. *Arabidopsis* trichome development has been extensively investigated, and several genes that control the process have been characterised. One gene essential for trichome initiation is *TRANSPARENT TESTA GLABRA1 (TTG1)*, and loss-of-function mutations in *TTG1* result in an almost complete absence of leaf trichomes. *TTG1* plays additional roles in numerous pathways in *Arabidopsis*, including root hair initiation, anthocyanin production and seed coat mucilage production. In order to isolate genes required for fibre initiation in cotton, functional homologues of *Arabidopsis TTG1* in cotton have been sought.

Four putative homologues of *Arabidopsis TTG1* have previously been isolated in this laboratory by RT-PCR of mRNA prepared from cotton fibres, and are termed GhTTG1-4. Sequence comparisons between the four cotton deduced proteins and *Arabidopsis TTG1* showed that they form two groups, with GhTTG1 and GhTTG3 being closely related to each other (87% identical and 93% similar) and to *TTG1* (79% and 80% amino acid identity respectively). GhTTG2 and GhTTG4 formed the second group, with 95% amino acid identity to each other and lower (approximately 62%) identity to *TTG1*. An analysis of the genomic origins of the *GhTTG* genes demonstrated that each is derived from the same ancestral diploid genome.

Cross-species complementation experiments were performed to test for functional homology of these cotton *TTG1*-like genes to *AtTTG1*, by introducing the cotton genes into *Arabidopsis ttg1-1* mutants via *Agrobacterium*-mediated transformation. This experiment

showed that two of the four genes, *GhTTG1* and *GhTTG3* were able to restore trichome initiation in the *Arabidopsis* mutant plants, and a further investigation of *GhTTG3* transgenic plants demonstrated complementation of the full range of *ttg1* mutant phenotypes.

An analysis of the temporal and spatial expression of the *GhTTG* genes in cotton is also described. It was shown that each of the genes is expressed ubiquitously throughout the cotton plant, in common with many plant WD-repeat genes. A closer examination of transcript abundance in the developing cotton ovule utilising *in situ* hybridisation revealed predominant expression of *GhTTG1/GhTTG3* in the epidermal cells destined to become cotton fibres.

A yeast two-hybrid assay was utilised to identify transcription factors that may interact with GhTTG3 during fibre development. This experiment identified three cotton fibre cDNAs encoding putative interacting proteins, including one with a similar secondary structure to several TTG1-interacting proteins in *Arabidopsis*, raising the possibility of similar regulatory complexes controlling trichome initiation in *Arabidopsis* and cotton.