



THE UNIVERSITY  

---

of ADELAIDE

# A Magnetic Spectrometer Analysis Method for Ultra High Energy Cosmic Ray Data

Benjamin James Whelan

B.Sc. (Honours), Physics

A thesis submitted to the University of Adelaide in fulfilment of the  
requirements for the degree of Doctor of Philosophy.

School of Chemistry & Physics

Department of Physics

November 2011

# Contents

<b>Abstract</b>	<b>i</b>
<b>Declaration</b>	<b>ii</b>
<b>Acknowledgements</b>	<b>iii</b>
<b>1 Introduction to Cosmic Rays and Extensive Air Showers</b>	<b>1</b>
1.1 The Discovery of Cosmic Rays and Extensive Air Showers . . . . .	1
1.2 Extensive Air Showers and Detection Techniques . . . . .	3
1.2.1 Formation and Morphology of Extensive Air Showers . . . . .	4
1.2.1.1 Heitler’s Model of Electromagnetic Cascades . . . . .	4
1.2.1.2 Hadronic Air Showers . . . . .	6
1.2.1.3 Atmospheric Fluorescence and Čerenkov Light Production . . . . .	11
1.2.2 Extensive Air Shower Detection . . . . .	13
1.2.2.1 Ground Arrays . . . . .	13
1.2.2.2 Fluorescence Detection . . . . .	16
1.2.2.3 ‘Hybrid’ Detection . . . . .	19
<b>2 Experimental Observations of the Cosmic Ray Flux</b>	<b>22</b>
2.1 The Energy Spectrum of Cosmic Rays . . . . .	22
2.2 Composition of Cosmic Rays . . . . .	28
2.3 Cosmic Ray Sources and Anisotropies . . . . .	30
2.3.1 Cosmic Ray Sources . . . . .	31
2.3.2 Recent Anisotropy Results at EeV Energies . . . . .	33
2.4 Interpretation of Energy Spectrum, Composition and Anisotropy Results . . . . .	37
<b>3 Ultra High Energy Cosmic Ray Detectors</b>	<b>39</b>
3.1 Previous UHECR Detectors . . . . .	39
3.1.1 Volcano Ranch . . . . .	39
3.1.2 Haverah Park . . . . .	40
3.1.3 SUGAR . . . . .	42

3.1.4	Yakutsk . . . . .	43
3.1.5	Fly’s Eye . . . . .	44
3.1.6	AGASA . . . . .	45
3.1.7	HiRes . . . . .	48
3.2	The Pierre Auger Observatory . . . . .	49
3.2.1	Communications and CDAS . . . . .	50
3.2.2	Surface Detector . . . . .	50
3.2.2.1	Station Design . . . . .	51
3.2.2.2	Calibration . . . . .	52
3.2.2.3	Trigger System . . . . .	53
3.2.2.4	Event Reconstruction and Performance . . . . .	56
3.2.3	Fluorescence Detector . . . . .	58
3.2.3.1	Telescope Design . . . . .	58
3.2.3.2	Electronics and Triggering . . . . .	59
3.2.3.3	Calibration and Atmospheric Monitoring . . . . .	60
3.2.3.4	Performance . . . . .	61
3.2.4	Hybrid Operation . . . . .	63
3.2.5	Auger South Enhancements . . . . .	63
3.2.5.1	AMIGA . . . . .	64
3.2.5.2	HEAT . . . . .	65
3.2.5.3	AERA . . . . .	65
3.2.6	Auger North . . . . .	66
<b>4</b>	<b>Cosmic Magnetic Fields and Cosmic Rays</b>	<b>70</b>
4.1	Cosmic Rays in Magnetic Fields . . . . .	70
4.2	Galactic Magnetic Fields . . . . .	73
4.2.1	Measurement Methods . . . . .	73
4.2.2	Magnetic Field Models . . . . .	74
4.2.2.1	Spiral Field . . . . .	75
4.2.2.2	Halo Fields . . . . .	76
4.2.2.3	Turbulent Fields . . . . .	78
<b>5</b>	<b>Numerical Simulations of Particle Propagation</b>	<b>80</b>
5.1	Numerical Integration . . . . .	80
5.1.1	Cosmic Ray Back-tracking . . . . .	83
5.2	Step Size and Accuracy . . . . .	84
5.3	Calculation of the Turbulent Field . . . . .	85
5.3.1	Choice of Minimum Turbulent Scale Size . . . . .	86

<b>6</b>	<b>Energy Ordering in UHECR Arrival Directions</b>	<b>92</b>
6.1	UHECR Energy Ordering and Analyses . . . . .	93
6.2	Introduction to the Magnetic Spectrometer Method . . . . .	99
6.3	Incorporating Experimental Uncertainties Into Source Position Reconstruction . . . . .	104
6.4	Application to Cosmic Ray Data Sets . . . . .	109
<b>7</b>	<b>Magnetic Spectrometer Analysis of Simulated UHECR Data</b>	<b>112</b>
7.1	Generation of Simulated Data Sets . . . . .	112
7.2	Analysis of Signal Events Only . . . . .	115
7.3	Analysis of Signal and Background Events . . . . .	120
	7.3.1 Peak-Finding Algorithm . . . . .	121
	7.3.2 Results of Magnetic Spectrometer Analysis . . . . .	122
	7.3.3 Estimation of Magnetic Fields . . . . .	126
7.4	Varying Relative Source Brightness . . . . .	130
7.5	Mixed Composition Source Flux . . . . .	133
<b>8</b>	<b>Analysis of Pierre Auger Observatory Data</b>	<b>136</b>
8.1	Analysis of Full Data Set . . . . .	136
8.2	Analysis of Epochs . . . . .	142
8.3	Interpretation of Results . . . . .	143
8.4	Conclusions and Future Prospects . . . . .	146
	<b>References</b>	<b>149</b>

# Abstract

Since their discovery at the beginning of the 20<sup>th</sup> Century, cosmic rays have been the subject of much research. However, fundamental questions, such as the origin of the most energetic of these particles, await an answer. The unambiguous identification of cosmic ray sources is made difficult by the deflections that these charged particles suffer in their propagation through cosmic magnetic fields. This problem is compounded by the extremely low flux of the highest energy cosmic rays at Earth, the arrival directions of which may be expected to retain information about their point of origin.

However, the advent of ultra high energy cosmic ray detectors possessing collecting areas of thousands of square kilometres, such as the Pierre Auger Observatory, means that the number of cosmic rays detected at the highest energies may be sufficient to enable directional cosmic ray astronomy in the near future. The ‘magnetic spectrometer analysis’, described in this thesis, is designed for the analysis of ultra high energy cosmic ray data sets. The analysis is designed to identify energy ordering in the arrival directions of a cosmic ray data set, and reconstruct the source directions of such events.

A brief history of the discovery of cosmic rays is presented in Chapter 1, along with an introduction to the physics of extensive air showers and methods of detecting them. The current knowledge of the properties of the cosmic ray flux at ultra high energies is reviewed in Chapter 2, and a summary of ultra high energy cosmic ray detectors, both past and present, is presented in Chapter 3.

The propagation of cosmic rays through magnetic fields, and the methods of measuring those fields, is briefly reviewed in Chapter 4. In addition, the Galactic magnetic field models that are used for the production of simulated cosmic ray data sets are described there.

Numerical integration is employed in the generation of the simulated data sets, and the method of doing so, as well as considerations that must be made for such simulations, is described in Chapter 5.

The magnetic spectrometer analysis method is introduced in Chapter 6 in addition to a discussion of related analyses found in the literature. The results of applying the magnetic spectrometer analysis to simulated cosmic ray data sets are presented in Chapter 7.

Finally, the results of an application of the magnetic spectrometer analysis to real data collected by the Pierre Auger Observatory are described in Chapter 8, followed by a discussion of those results and concluding remarks.

# Declaration of Originality

I, Benjamin James Whelan, certify that this work contains no material which has been accepted for the award of any other degree or diploma in any university or other tertiary institution and, to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference has been made in the text.

I give consent to this copy of my thesis, when deposited in the University Library, being made available for loan and photocopying, subject to the provisions of the Copyright Act 1968.

I also give permission for the digital version of my thesis to be made available on the web, via the University's digital research repository, the Library catalogue and also through web search engines, unless permission has been granted by the University to restrict access for a period of time.

**Signed:** .....

**Date:** .....

# Acknowledgements

I would like to thank my supervisors, Roger Clay and Bruce Dawson, for the help and guidance they have offered me over the course of my Ph.D. They have taught me much, and I am grateful for the enthusiasm and interest toward my research they have shown during my studies. In addition, their eternal patience and calming influence was invaluable on the many occasions when I felt things were all getting to be too much.

I would also like to thank the other members of the High Energy Astrophysics research group at the University of Adelaide, both past and present. They have made my time as a Ph.D. student an interesting and enlightening one, and have been ever willing to offer computing expertise or help tackle a difficult problem. Particular thanks go to my office mate, Kerri Budd, who brings a cheerful mood to the office every morning. She has always been ready to talk over any issues I have had and provide useful input. In addition, her willingness to offer a fresh set of eyes whenever required was much appreciated.

The Pierre Auger Collaboration ([www.auger.org/admin/Collaborators/author\\_list\\_alphabetical.html](http://www.auger.org/admin/Collaborators/author_list_alphabetical.html)) has been an exciting group of people to work with, and I thank the collaboration members for the input and ideas that they have offered towards this work. The opportunity to work with a cutting edge detector, and be exposed to so many different areas of the field, has given me an excellent start to cosmic ray research.

I was fortunate enough to be awarded the Ferry Postgraduate Scholarship, for which I am extremely grateful. Computing facilities provided by eResearch SA were used extensively throughout my research, which allowed me to generate much of the data presented in this thesis.

My family and friends have been a constant source of support over the course of my studies. In particular, I extend thanks to my parents, Marg and Ken, and my sister, Kate. They have always encouraged me to do whatever makes me happy, and I couldn't ask for more.

Last, but most certainly not least, is my wife, Alex. An unwavering source of love, she has kept me going through even the most stressful periods of my Ph.D. It's an incredible feeling to have her believe in me, and I can't imagine having gotten this far without her.